arXiv:0905.3117v1 [math.QA] 19 May 2009

CENTERS OF GRADED FUSION CATEGORIES

SHLOMO GELAKI, DEEPAK NAIDU, AND DMITRI NIKSHYCH

Abstract. Let $\mathcal{C}$ be a fusion category faithfully graded by a finite group G and let $\mathcal{D}$ be the trivial component of this grading. The center $\mathcal{Z}(\mathcal{C})$ of $\mathcal{C}$ is shown to be canonically equivalent to a G-equivariantization of the relative center $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. We use this result to obtain a criterion for $\mathcal{C}$ to be group-theoretical and apply it to Tambara-Yamagami fusion categories. We also find several new series of modular categories by analyzing the centers of Tambara-Yamagami categories. Finally, we prove a general result about existence of zeroes in S-matrices of weakly integral modular categories.

Contents

1. Introduction 2
2. Preliminaries 3
3. The center of a graded fusion category 10
4. The centers of Tambara-Yamagami categories 14
5. Examples of modular categories arising from quadratic forms 19
6. Appendix: Zeroes in S-matrices 24
References 26

1. Introduction

Throughout the paper we work over an algebraically closed field k of characteristic 0. All categories considered in this paper are finite, Abelian, semisimple, and k-linear. We freely use the language and basic theory of fusion categories, module categories over them, braided categories, and Frobenius-Perron dimensions [BK, O, ENO1].

Let G be a finite group. A fusion category $\mathcal{C}$ is *G-graded* if there is a decomposition

$$\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g$$

of $\mathcal{C}$ into a direct sum of full Abelian subcategories such that the tensor product of $\mathcal{C}$ maps $\mathcal{C}_g \times \mathcal{C}_h$ to $\mathcal{C}_{gh}$, for all $g, h \in G$. A *G-extension* of a fusion category $\mathcal{D}$ is a G-graded fusion category $\mathcal{C}$ whose trivial component $\mathcal{C}_e$, where e is the identity of G, is equivalent to $\mathcal{D}$.

Gradings and extensions play an important role in the study and classification of fusion categories. E.g., *nilpotent* fusion categories (i.e., those categories that can be obtained from the trivial category by a sequence of groups extensions) were studied in [GN]. It was proved in [ENO1] that every fusion category of prime power dimension is nilpotent. Group-theoretical properties of such categories were studied in [DGNO]. Recently, fusion categories of dimension $p^n q^m$, where p, q are primes, were shown to be Morita equivalent to nilpotent categories [ENO3].

The main goal of this paper is to describe the center $\mathcal{Z}(\mathcal{C})$ of a G-graded fusion category $\mathcal{C}$ in terms of its trivial component $\mathcal{D}$ (Theorem 3.5) and apply this description to the study of structural properties of $\mathcal{C}$ and construction of new examples of modular categories.

The organization of the paper is as follows. In Section 2 we recall some basic notions, results, and examples of fusion categories, notably the notions of the relative center of a bimodule category [Ma], group action on a fusion category and crossed product [Ta2], equivariantization and de-equivariantization theory [AG, Br, G, Ki, Mu1, DGNO], and braided G-crossed fusion categories [Tu1, Tu2].

In Section 3 we study the center $\mathcal{Z}(\mathcal{C})$ of a G-graded fusion category $\mathcal{C}$. We show that if $\mathcal{D}$ is the trivial component of $\mathcal{C}$, then the relative center $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ has a canonical structure of a braided G-crossed category and there is an equivalence of braided fusion categories $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})^G \cong \mathcal{Z}(\mathcal{C})$ (Theorem 3.5). Thus, the structure of $\mathcal{Z}(\mathcal{C})$ can be understood in terms of a smaller and more transparent category $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. In particular, there is a canonical braided action of G on $\mathcal{Z}(\mathcal{D})$[1]. In Corollary 3.10 we use this action to prove that $\mathcal{C}$ is group-theoretical if and only if $\mathcal{Z}(\mathcal{D})$ contains a G-stable Lagrangian subcategory. As an illustration, we describe the center of a crossed product fusion category $\mathcal{C} = \mathcal{D} \rtimes G$.

We apply the above results in Section 4 to the study of Tambara-Yamagami categories [TY]. We obtain a convenient description of the centers of such categories as equivariantizations and compute their modular data, i.e., S- and T-matrices. This computation was previously done by Izumi in [I] using different techniques. We establish a criterion for a Tambara-Yamagami category to be group-theoretical (Theorem 4.6). We also extend the construction of non group-theoretical semisimple Hopf algebras from Tambara-Yamagami categories given in [Ni].

[1]This action is studied in detail in [ENO3].

In Section 5 we construct a series of new modular categories as factors of the centers of Tambara-Yamagami categories. Namely, one associates a pair of such categories $\mathcal{E}(q, \pm)$ with any non-degenerate quadratic form q on an Abelian group A of odd order. The categories $\mathcal{E}(q, \pm)$ have dimension $4|A|$. They are group-theoretical if and only if A contains a Lagrangian subgroup with respect to q. We compute the S- and T- matrices of $\mathcal{E}(q, \pm)$ and write down several small examples explicitly.

Section 6 is independent from the rest of the paper and contains a general result about existence of zeroes in S-matrices of weakly integral modular categories (Theorem 6.1). This is a categorical analogue of a classical result of Burnside in character theory.

Acknowledgments. We are grateful to P. Etingof, M. Müger, and V. Ostrik for useful discussions. Part of this work was done while the first author was on Sabbatical in the Departments of Mathematics at the University of New Hampshire and MIT; he is grateful for their warm hospitality. The research of S. Gelaki was partially supported by the Israel Science Foundation (grant No. 125/05). The research of D. Nikshych was supported by the NSA grant H98230-07-1-0081 and the NSF grant DMS-0800545.

2. Preliminaries

Below we recall several constructions and results used in the sequel.

2.1. **Dual fusion categories and Morita equivalence.** Let $\mathcal{C}$ be a fusion category and let $\mathcal{M}$ be an indecomposable right $\mathcal{C}$-module category $\mathcal{M}$. The category $\mathcal{C}^*_{\mathcal{M}}$ of $\mathcal{C}$-module endofunctors of $\mathcal{M}$ is a fusion category, called the dual of $\mathcal{C}$ with respect to $\mathcal{M}$ (see [ENO1, O]).

Following [Mu3], we say that two fusion categories $\mathcal{C}$ and $\mathcal{D}$ are *Morita equivalent* if $\mathcal{D}$ is equivalent to $\mathcal{C}^*_{\mathcal{M}}$, for some indecomposable right $\mathcal{C}$-module category $\mathcal{M}$. A fusion category is said to be *pointed* if all its simple objects are invertible (any such category is equivalent to the category Vec_G^ω of vector spaces graded by a finite group G with the associativity constraint given by a 3-cocylce $\omega \in Z^3(G, k^\times)$). A fusion category is called *group-theoretical* if it is Morita equivalent to a pointed fusion category. See [O, ENO1, Ni] for details of the theory of group-theoretical categories.

2.2. **The center of a bimodule category and the relative center of a fusion category.** Let $\mathcal{C}$ be a fusion category with unit object $\mathbf{1}$ and associativity constraint $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \xrightarrow{\sim} X \otimes (Y \otimes Z)$ and let $\mathcal{M}$ be a $\mathcal{C}$-bimodule category.

Definition 2.1. The *center* of $\mathcal{M}$ is the category $\mathcal{Z}_{\mathcal{C}}(\mathcal{M})$ of $\mathcal{C}$-bimodule functors from $\mathcal{C}$ to $\mathcal{M}$.

Explicitly, the objects of $\mathcal{Z}_{\mathcal{C}}(\mathcal{M})$ are pairs (M, γ), where M is an object of $\mathcal{M}$ and

$$\gamma = \{\gamma_X : X \otimes M \xrightarrow{\sim} M \otimes X\}_{X \in \mathcal{C}} \tag{1}$$

is a natural family of isomorphisms making the following diagram commutative:
(2)



where α's denote the associativity constraints in $\mathcal{M}$.

Indeed, a $\mathcal{C}$-bimodule functor $F : \mathcal{C} \to \mathcal{M}$ is completely determined by the pair $(F(\mathbf{1}), \{\gamma_X\}_{X\in\mathcal{C}})$, where $\gamma = \{\gamma_X\}_{X\in\mathcal{C}}$ is the collection of isomorphisms

$$\gamma_X : X \otimes F(\mathbf{1}) \xrightarrow{\sim} F(X) \xrightarrow{\sim} F(\mathbf{1}) \otimes X$$

coming from the $\mathcal{C}$-bimodule structure on F.

We will call the natural family of isomorphisms (1) the *central structure* of an object $X \in \mathcal{Z}_{\mathcal{C}}(\mathcal{M})$.

Remark 2.2. (i) The definition of the center of a bimodule category is parallel to that of the center of a bimodule over a ring.

(ii) We will often suppress the central structure while working with objects of $\mathcal{Z}_{\mathcal{C}}(\mathcal{M})$ and refer to (M, γ) simply as M.

(iii) $\mathcal{Z}_{\mathcal{C}}(\mathcal{M})$ is a semisimple Abelian category. It has an obvious canonical structure of a $\mathcal{Z}(\mathcal{C})$-module category, where $\mathcal{Z}(\mathcal{C})$ is the center of $\mathcal{C}$ (see e.g., [K, Section XIII.4] for the definition of $\mathcal{Z}(\mathcal{C})$).

Here is an important special case of the above construction. Let $\mathcal{C}$ be a fusion category and let $\mathcal{D} \subset \mathcal{C}$ be a fusion subcategory. Then $\mathcal{C}$ is a $\mathcal{D}$-bimodule category. We will call $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ the *relative center* of $\mathcal{C}$.

Remark 2.3. The aforementioned construction of relative center is a special case of a more general construction considered by Majid in [Ma] (see Definition 3.2 and Theorem 3.3 of [Ma]).

It is easy to see that $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ is a tensor category with tensor product defined as follows. If (X, γ) and (X', γ') are objects in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ then

$$(X, \gamma) \otimes (X', \gamma') := (X \otimes X', \tilde{\gamma}),$$

where $\tilde{\gamma}_V : V \otimes (X \otimes X') \xrightarrow{\sim} (X \otimes X') \otimes V$, $V \in \mathcal{D}$, is defined by the following diagram:

(3)



The unit object of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ is $(\mathbf{1}, \mathrm{id})$. The dual of (X, γ) is $(X^*, \overline{\gamma})$, where $\overline{\gamma}_V := (\gamma_{*V})^*$.

Remark 2.4. Let $\mathcal{C}$ and $\mathcal{D}$ be as above.

(i) $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ is dual to the fusion category $\mathcal{D} \boxtimes \mathcal{C}^{\mathrm{rev}}$ (where $\mathcal{C}^{\mathrm{rev}}$ is the fusion category obtained from $\mathcal{C}$ by reversing the tensor product and $\boxtimes$ is Deligne's tensor product of fusion categories) with respect to its module category $\mathcal{C}$, where $\mathcal{D}$ and $\mathcal{C}^{\mathrm{rev}}$ act on $\mathcal{C}$ via the right and left multiplication respectively. In particular, $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ is a fusion category.

(ii) $\mathrm{FPdim}(\mathcal{Z}_{\mathcal{D}}(\mathcal{C})) = \mathrm{FPdim}(\mathcal{C})\,\mathrm{FPdim}(\mathcal{D})$, where FPdim denotes the Frobenius-Perron dimension of a category.

(iii) $\mathcal{Z}_{\mathcal{C}}(\mathcal{C})$ coincides with the center $\mathcal{Z}(\mathcal{C})$ of $\mathcal{C}$. This category has a canonical braiding given by

$$c_{(X,\gamma),\,(X',\gamma')} = \gamma_{X'} : (X,\,\gamma) \otimes (X',\,\gamma') \xrightarrow{\sim} (X',\,\gamma') \otimes (X,\,\gamma). \tag{4}$$

(iv) There is an obvious forgetful tensor functor:

$$\mathcal{Z}(\mathcal{C}) \mapsto \mathcal{Z}_{\mathcal{D}}(\mathcal{C}) : (X,\,\gamma) \mapsto (X,\,\gamma|_{\mathcal{D}}). \tag{5}$$

2.3. **Centralizers in braided fusion categories.** Let $\mathcal{C}$ be a braided fusion category with braiding c. Two objects X and Y of $\mathcal{C}$ are said to *centralize* each other [Mu2] if $c_{Y,X}c_{X,Y} = \mathrm{id}_{X\otimes Y}$.

For any fusion subcategory $\mathcal{D} \subseteq \mathcal{C}$ its *centralizer* $\mathcal{D}'$ is the full fusion subcategory of $\mathcal{C}$ consisting of all objects $X \in \mathcal{C}$ which centralizes every object in $\mathcal{D}$. The category $\mathcal{C}$ is said to be *non-degenerate* if $\mathcal{C}' = \mathrm{Vec}$. In this case one has $\mathcal{D}'' = \mathcal{D}$ [Mu2]. If $\mathcal{C}$ is a pre-modular category, i.e., has a spherical structure, then it is non-degenerate if and only if it is modular.

A braided fusion category $\mathcal{E}$ is called *Tannakian* if it is equivalent to the representation category $\mathrm{Rep}(G)$ of a finite group G as a braided fusion category. Here $\mathrm{Rep}(G)$ is considered with its standard symmetric braiding. The group G is defined by $\mathcal{E}$ up to an isomorphism [D].

A fusion subcategory $\mathcal{L}$ of a braided fusion category is called *Lagrangian* if it is Tannakian and $\mathcal{L} = \mathcal{L}'$.

Theorem 2.5 ([DGNO])**.** *A fusion category $\mathcal{C}$ is group-theoretical if and only if $\mathcal{Z}(\mathcal{C})$ contains a Lagrangian subcategory.*

2.4. **Group actions on fusion categories and equivariantization.** Let G be a finite group, and let $\underline{G}$ denote the monoidal category whose objects are elements of G, morphisms are identities, and the tensor product is given by the multiplication in G. Recall that an action of G on a fusion category $\mathcal{C}$ is a monoidal functor $\underline{G} \to \mathrm{Aut}_{\otimes}(\mathcal{C}) : g \mapsto T_g$. For any $g, h \in G$, let $\gamma_{g,h}$ be the isomorphism $T_g \circ T_h \simeq T_{gh}$ that defines the monoidal structure on the functor $\underline{G} \to \mathrm{Aut}_{\otimes}(\mathcal{C})$.

Definition 2.6. A *G-equivariant object* in $\mathcal{C}$ is a pair $(X, \{u_g\}_{g\in G})$ consisting of an object X of $\mathcal{C}$ together with a collection of isomorphisms $u_g : T_g(X) \simeq X$, $g \in G$, such that the diagram

$$\begin{array}{ccc} T_g(T_h(X)) & \xrightarrow{T_g(u_h)} & T_g(X) \\ {\scriptstyle \gamma_{g,h}(X)}\downarrow & & \downarrow{\scriptstyle u_g} \\ T_{gh}(X) & \xrightarrow{u_{gh}} & X \end{array}$$

commutes for all $g, h \in G$. One defines morphisms of equivariant objects to be morphisms in $\mathcal{C}$ commuting with $u_g,\ g \in G$.

Equivariant objects in $\mathcal{C}$ form a fusion category, called the *equivariantization* of $\mathcal{C}$ and denoted by $\mathcal{C}^G$, see [Ta2, AG, G]. One has $\mathrm{FPdim}(\mathcal{C}^G) = |G|\,\mathrm{FPdim}(\mathcal{C})$.

There is another fusion category that comes from an action of G on $\mathcal{C}$. It is the *crossed product* category $\mathcal{C} \rtimes G$ defined as follows, see [Ta2, Ni]. As an Abelian category, $\mathcal{C} \rtimes G := \mathcal{C} \boxtimes \mathrm{Vec}_G$, where Vec_G denotes the fusion category of G-graded vector spaces. The tensor product in $\mathcal{C} \rtimes G$ is given by

$$(X \boxtimes g) \otimes (Y \boxtimes h) := (X \otimes T_g(Y)) \boxtimes gh, \qquad X, Y \in \mathcal{C}, \quad g, h \in G. \tag{6}$$

The unit object is $\mathbf{1} \boxtimes e$ and the associativity and unit constraints come from those of $\mathcal{C}$. Clearly, $\mathcal{C} \rtimes G$ is faithfully G-graded with the trivial component $\mathcal{C}$.

It was explained in [Ni] that $\mathcal{C}$ is a right $\mathcal{C} \rtimes G$-module category via

$$Y \otimes (X \boxtimes g) := T_{g^{-1}}(Y \otimes X)$$

and the corresponding dual category $(\mathcal{C} \rtimes G)^*_{\mathcal{C}}$ is equivalent to $\mathcal{C}^G$. It follows from [Mu3] that there is an equivalence of braided fusion categories

$$\mathcal{Z}(\mathcal{C} \rtimes G) \cong \mathcal{Z}(\mathcal{C}^G).$$

Let G be a finite group. For any conjugacy class K of G fix a representative $a_K \in K$. Let G_K denote the centralizer of a_K in G.

Proposition 2.7. *Let $\mathcal{C} = \bigoplus_{g \in G} \mathcal{C}_g$ be a G-graded fusion category with an action $g \mapsto T_g$ of G on $\mathcal{C}$ such that T_g carries $\mathcal{C}_h$ to $\mathcal{C}_{ghg^{-1}}$. Let $H := \{g \in G \mid \mathcal{C}_g \neq 0\}$. There is a bijection between the set of isomorphism classes of simple objects of $\mathcal{C}^G$ and pairs (K, X), where $K \subset H$ is a conjugacy class of G and X is a simple G_K-equivariant object of $\mathcal{C}_{a_K}$.*

Proof. A simple G-equivariant object of $\mathcal{C}$ must be supported on a single conjugacy class K. Let $Y = \oplus_{g \in K} Y_g$ be such an object. Then Y_{a_K} is a simple G_K-equivariant object.

Conversely, given a G_K-equivariant object X in $\mathcal{C}_{a_K}$ let

$$Y = \bigoplus_h T_h(X),$$

where the summation is taken over the set of representatives of cosets of G_K in G. It is easy to see that Y acquires the structure of a simple G-equivariant object.

Clearly, the above constructions are inverses of each other. □

Remark 2.8. The Frobenius-Perron dimension of the simple object corresponding to a pair (K, X) in Proposition 2.7 is $|K|\,\mathrm{FPdim}(X)$.

2.5. **De-equivariantization of fusion categories.** Let $\mathcal{C}$ be a fusion category. Let $\mathcal{E} = \mathrm{Rep}(G)$ be a Tannakian category along with a braided tensor functor $\mathcal{E} \to \mathcal{Z}(\mathcal{C})$ such that the composition $\mathcal{E} \to \mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ (where the second arrow is the forgetful functor) is fully faithful. The following construction was introduced by Bruguières [Br] and Müger [Mu1]. Let $A := \mathrm{Fun}(G)$ be the algebra of functions on G. It is a commutative algebra in $\mathcal{E}$, hence, its image is a commutative algebra in $\mathcal{Z}(\mathcal{C})$. This fact allows to view the category $\mathcal{C}_G$ of A-modules in $\mathcal{C}$ as a fusion category, called *de-equivariantization* of $\mathcal{C}$. There is a canonical surjective tensor functor

$$F : \mathcal{C} \to \mathcal{C}_G : X \mapsto A \otimes X. \tag{7}$$

It was explained in [Mu1, DGNO] that the group G acts on $\mathcal{C}_G$ by tensor autoequivalences (this action comes from the action of G on A by right translations). Furthermore, there is a bijection between subcategories of $\mathcal{C}$ containing the image of $\mathcal{E} = \mathrm{Rep}(G)$ and G-stable subcategories of $\mathcal{C}_G$. This bijection preserves Tannakian subcategories.

The procedures of equivariantization and de-equivariantization are inverses of each other, i.e., there are canonical equivalences $(\mathcal{C}_G)^G \cong \mathcal{C}$ and $(\mathcal{C}^G)_G \cong \mathcal{C}$.

In particular, the above construction applies when $\mathcal{C}$ is a braided fusion category containing a Tannakian subcategory $\mathcal{E} = \mathrm{Rep}(G)$. In this case the braiding of $\mathcal{C}$ gives rise to an additional structure on the de-equivariantization functor (7). Namely, there is natural family of isomorphisms

$$X \otimes F(Y) \xrightarrow{\sim} F(Y) \otimes X, \qquad X \in \mathcal{C}_G,\, Y \in \mathcal{C}, \tag{8}$$

satisfying obvious compatibility conditions. In other words, F can be factored through a braided functor $\mathcal{C} \to \mathcal{Z}(\mathcal{C}_G)$, i.e., F is a *central* functor.

If $\mathcal{E} \subset \mathcal{C}'$ then $\mathcal{C}_G$ is a braided fusion category with the braiding inherited from that of $\mathcal{C}$. If $\mathcal{E} = \mathcal{C}'$, the category $\mathcal{C}_G$ is non-degenerate (in the presence of a spherical structure this category is called the *modularization* of $\mathcal{C}$ by $\mathcal{E}$ [Br, Mu1]).

Remark 2.9. The category $\mathcal{C}_G$ is not braided in general. However it does have an additional structure, namely it is a *braided G-crossed fusion category*. See Section 2.6 below for details.

2.6. **Braided G-crossed categories.** Let G be a finite group. Kirillov Jr. [Ki] and Müger [Mu4] found a description of all braided fusion categories $\mathcal{D}$ containing $\mathrm{Rep}(G)$. Namely, they showed that the datum of a braided fusion category $\mathcal{D}$ containing $\mathrm{Rep}(G)$ is equivalent to the datum of a braided G-crossed category $\mathcal{C}$, see Theorem 2.12. The notion of a braided G-crossed category is due to Turaev [Tu1, Tu2] and is recalled below.

Definition 2.10. A *braided G-crossed fusion category* is a fusion category $\mathcal{C}$ equipped with the following structures:

(i) a (not necessarily faithful) grading $\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g$,
(ii) an action $g \mapsto T_g$ of G on $\mathcal{C}$ such that $T_g(\mathcal{C}_h) \subset \mathcal{C}_{ghg^{-1}}$,
(iii) a natural collection of isomorphisms, called the *G-braiding*:

$$c_{X,Y} : X \otimes Y \simeq T_g(Y) \otimes X, \qquad X \in \mathcal{C}_g,\, g \in G \text{ and } Y \in \mathcal{C}. \tag{9}$$

Let $\gamma_{g,h} : T_g T_h \xrightarrow{\sim} T_{gh}$ denote the tensor structure of the functor $g \mapsto T_g$ and let μ_g denote the tensor structure of T_g.

The above structures are required to satisfy the following compatibility conditions:

(a) the diagram

(10)
$$\begin{array}{ccc}
T_g(X)\otimes T_g(Y) & \xrightarrow{c_{T_g(X),T_g(Y)}} & T_{ghg^{-1}}(T_g(Y))\otimes T_g(X) \\
{\scriptstyle (\mu_g)^{-1}_{X,Y}}\uparrow & & \downarrow{\scriptstyle (\gamma_{ghg^{-1},g})_Y\otimes \mathrm{id}_{T_g(X)}} \\
T_g(X\otimes Y) & & T_{gh}(Y)\otimes T_g(X) \\
{\scriptstyle T_g(c_{X,Y})}\downarrow & & \uparrow{\scriptstyle (\gamma_{g,h})_Y\otimes \mathrm{id}_{T_g(X)}} \\
T_g(T_h(Y)\otimes X) & \xrightarrow[(\mu_g)^{-1}_{T_g(Y),X}]{} & T_g(T_h(Y))\otimes T_g(X),
\end{array}$$

commutes for all $g, h \in G$ and objects $X \in \mathcal{C}_h$, $Y \in \mathcal{C}$,

(b) the diagram

(11)
$$\begin{array}{ccc}
 & (X\otimes Y)\otimes Z & \\
{\scriptstyle \alpha_{X,Y,Z}}\swarrow & & \searrow{\scriptstyle c_{X,Y}\otimes \mathrm{id}_Z} \\
X\otimes (Y\otimes Z) & & (T_g(Y)\otimes X)\otimes Z \\
{\scriptstyle c_{X,Y\otimes Z}}\downarrow & & \downarrow{\scriptstyle \alpha_{T_g(Y),X,Z}} \\
T_g(Y\otimes Z)\otimes X & & T_g(Y)\otimes (X\otimes Z) \\
{\scriptstyle (\mu_g)^{-1}_{Y,Z}\otimes \mathrm{id}_X}\downarrow & & \downarrow{\scriptstyle \mathrm{id}_{T_g(Y)}\otimes c_{X,Z}} \\
(T_g(Y)\otimes T_g(Z))\otimes X & \xrightarrow{\alpha_{T_g(Y),T_g(Z),X}} & T_g(Y)\otimes (T_g(Z)\otimes X)
\end{array}$$

commutes for all $g \in G$ and objects $X \in \mathcal{C}_g, Y, Z \in \mathcal{C}$, and

(c) the diagram

(12)
$$\begin{array}{ccc}
 & X\otimes (Y\otimes Z) & \\
{\scriptstyle \alpha_{X,Y,Z}}\nearrow & & \searrow{\scriptstyle \mathrm{id}_X\otimes c_{Y,Z}} \\
(X\otimes Y)\otimes Z & & X\otimes (T_h(Z)\otimes Y) \\
{\scriptstyle c^{-1}_{X\otimes Y,Z}}\uparrow & & \downarrow{\scriptstyle \alpha^{-1}_{X,T_h(Z),Y}} \\
T_{gh}(Z)\otimes (X\otimes Y) & & (X\otimes T_h(Z))\otimes Y \\
{\scriptstyle (\gamma_{g,h})_Z\otimes \mathrm{id}_{X\otimes Y}}\uparrow & & \downarrow{\scriptstyle c_{X,T_h(Z)}\otimes \mathrm{id}_Y} \\
T_gT_h(Z)\otimes (X\otimes Y) & \xrightarrow{\alpha^{-1}_{T_gT_h(Z),X,Y}} & (T_gT_h(Z)\otimes X)\otimes Y.
\end{array}$$

commutes for all $g, h \in G$ and objects $X \in \mathcal{C}_g$, $Y \in \mathcal{C}_h, Z \in \mathcal{C}$.

Remark 2.11. The trivial component $\mathcal{C}_e$ of a braided G-crossed fusion category $\mathcal{C}$ is a braided fusion category with the action of G by braided autoequivalences. This can be seen by taking X, $Y \in \mathcal{C}_e$ in diagrams (10) – (12).

Theorem 2.12 ([Ki, Mu4])**.** *The equivariantization and de-equivariantization constructions establish a bijection between the set of equivalence classes of G-crossed*

braided fusion categories and the set of equivalence classes of braided fusion categories containing $\mathrm{Rep}(G)$ *as a symmetric fusion subcategory.*

We shall now sketch the proof of this theorem. An alternative approach is given in [DGNO].

Suppose $\mathcal{C}$ is a braided G-crossed fusion category. We define a braiding $\tilde{c}$ on its equivariantization $\mathcal{C}^G$ as follows.

Let $(X, \{u_g\}_{g\in G})$ and $(Y, \{v_g\}_{g\in G})$ be objects in $\mathcal{C}^G$. Let $X = \oplus_{g\in G} X_g$ be a decomposition of X with respect to the grading of $\mathcal{C}$. Define an isomorphism

(13)
$$\tilde{c}_{X,Y} : X\otimes Y = \bigoplus_{g\in G} X_g\otimes Y \xrightarrow{\oplus c_{X_g,Y}} \bigoplus_{g\in G} T_g(Y)\otimes X_g \xrightarrow{\oplus v_g\otimes \mathrm{id}_{X_g}} \bigoplus_{g\in G} Y\otimes X_g = Y\otimes X,$$

It follows from condition (a) of Definition 2.10 that $\tilde{c}_{X,Y}$ respects the equivariant structures, i.e., it is an isomorphism in $\mathcal{C}^G$. Its naturality is clear. The fact that $\tilde{c}$ is a braiding on $\mathcal{C}^G$ (i.e., the hexagon axioms) follows from the commutativity of diagrams (11) and (12). It is easy to check that $\tilde{c}$ restricts to the standard braiding on $\mathrm{Rep}(G) = \mathrm{Vec}^G \subset \mathcal{C}^G$. Hence, $\mathcal{C}^G$ contains a Tannakian subcategory $\mathrm{Rep}(G)$.

Conversely, let $\mathcal{C}$ be a braided fusion category with braiding c containing a Tannakian subcategory $\mathrm{Rep}(G)$. The restriction of the de-equivariantization functor F from (7) on $\mathrm{Rep}(G)$ is isomorphic to the fiber functor $\mathrm{Rep}(G) \to \mathrm{Vec}$. Hence for any object X in $\mathcal{C}_G$ and any object V in $\mathrm{Rep}(G)$ we have an automorphism of $F(V)\otimes X$ defined as the composition

$$F(V)\otimes X \xrightarrow{\sim} X\otimes F(V) \xrightarrow{\sim} F(V)\otimes X, \tag{14}$$

where the first isomorphism comes from the fact that $F(V) \in \mathrm{Vec}$ and the second one is (8).

When X is simple we have an isomorphism $\mathrm{Aut}_{\mathcal{C}}(F(V)\otimes X) \cong \mathrm{Aut}_{\mathrm{Vec}}(F(V))$, hence we obtain a tensor automorphism i_X of $F|_{\mathrm{Rep}(G)}$. Since $\mathrm{Aut}_{\otimes}(F|_{\mathrm{Rep}(G)}) \cong G$ we have an assignment $X \mapsto i_X \in G$. The hexagon axiom of braiding implies that this assignment is multiplicative, i.e., that $i_Z = i_X i_Y$ for any simple object Z contained in $X\otimes Y$. Thus, it defines a G-grading on $\mathcal{C}$:

$$\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g, \text{ where } \mathcal{O}(\mathcal{C}_g) = \{X \in \mathcal{O}(\mathcal{C}) \mid i_X = g\}. \tag{15}$$

It is straightforward to check that $i_{T_g(X)} = ghg^{-1}$ whenever $i_X = h$.

Finally, to construct a G-crossed braiding on $\mathcal{C}$ observe that $\mathcal{C}$ and $\mathcal{C}^{\mathrm{rev}}$ are embedded into the crossed product category $\mathcal{C}\rtimes G = (\mathcal{C}^G)^*_{\mathcal{C}}$ as subcategories $\mathcal{C}_{\mathrm{left}}$ and $\mathcal{C}_{\mathrm{right}}$ consisting, respectively, of functors of left and right multiplications by objects of $\mathcal{C}$. Clearly, there is a natural family of isomorphisms

$$X\otimes Y \xrightarrow{\sim} Y\otimes X, \qquad X\in\mathcal{C}_{\mathrm{left}}, Y\in\mathcal{C}_{\mathrm{right}}, \tag{16}$$

satisfying obvious compatibility conditions. Note that $\mathcal{C}_{\mathrm{left}}$ is identified with the diagonal subcategory of $\mathcal{C}\rtimes G$ spanned by objects $X\boxtimes g$, $X\in\mathcal{C}_g$, $g\in G$, and $\mathcal{C}_{\mathrm{right}}$ is identified with the trivial component subcategory $\mathcal{C}\boxtimes e$. Using (6) we conclude that isomorphisms (16) give rise to a G-crossed braiding on $\mathcal{C}$.

One can check that the two above constructions (from braided fusion categories containing $\mathrm{Rep}(G)$ to braided G-crossed categories and vice versa) are inverses of each other, see [Ki, Mu4, DGNO] for details.

Remark 2.13. Let $\mathcal{C} = \oplus_{g\in G}\mathcal{C}_g$ be a braided G-crossed fusion category. It was shown in [DGNO] that the braided category $\mathcal{C}^G$ is non-degenerate if and only if $\mathcal{C}_e$ is non-degenerate and the G-grading of $\mathcal{C}$ is faithful.

3. The center of a graded fusion category

Let G be a finite group and let $\mathcal{D}$ be a fusion category. Throughout this section $\mathcal{C}$ will denote a fusion category with a faithful G-grading, whose trivial component is $\mathcal{D}$, i.e., $\mathcal{C}$ is a G-extension of $\mathcal{D}$:

$$\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g, \qquad \mathcal{C}_e = \mathcal{D}. \tag{17}$$

In what follows we consider only *faithful* gradings, i.e., such that $\mathcal{C}_g \neq 0$, for all $g \in G$. An object of $\mathcal{C}$ contained in $\mathcal{C}_g$ will be called *homogeneous* of degree g.

Our goal is to describe the center $\mathcal{Z}(\mathcal{C})$ as an equivariantization of the relative center $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ defined in Section 2.2.

3.1. **The relative center $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ as a braided G-crossed category.** Let us define a canonical braided G-crossed category structure on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$.

First of all, there is an obvious faithful G-grading on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$:

$$\mathcal{Z}_{\mathcal{D}}(\mathcal{C}) = \bigoplus_{g\in G} \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g). \tag{18}$$

Indeed, it is clear that for every simple object X of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ the forgetful image of X in $\mathcal{C}$ must be homogeneous.

Next, let us define the action of G on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. Take $g, h \in G$.

Let $\mathrm{Fun}_{\mathcal{D}\boxtimes\mathcal{D}^{\mathrm{rev}}}(\mathcal{C}_g, \mathcal{C}_h)$ denote the category of $\mathcal{D}$-bimodule functors from $\mathcal{C}_g$ to $\mathcal{C}_h$. Clearly, it is a $\mathcal{Z}(\mathcal{D})$-bimodule category.

Proposition 3.1. *Let $g, h \in G$. The functors*

$$L_{g,h} : \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_h) \xrightarrow{\sim} \mathrm{Fun}_{\mathcal{D}\boxtimes\mathcal{D}^{rev}}(\mathcal{C}_g, \mathcal{C}_{hg}) : Z \mapsto Z\otimes ?, \tag{19}$$

$$R_{g,h} : \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_h) \xrightarrow{\sim} \mathrm{Fun}_{\mathcal{D}\boxtimes\mathcal{D}^{rev}}(\mathcal{C}_g, \mathcal{C}_{gh}) : Z \mapsto ? \otimes Z. \tag{20}$$

are equivalences of $\mathcal{Z}(\mathcal{D})$-bimodule categories.

Proof. We prove that (19) is an equivalence. Let $\mathrm{Fun}_{\mathcal{D}}(\mathcal{C}_g, \mathcal{C}_{hg})$ be the category of right $\mathcal{D}$-module functors from $\mathcal{C}_g$ to $\mathcal{C}_{hg}$. It suffices to prove that

$$M_{g,h} : \mathcal{C}_h \to \mathrm{Fun}_{\mathcal{D}}(\mathcal{C}_g, \mathcal{C}_{hg}) : X \mapsto X\otimes ? \tag{21}$$

is an equivalence. Indeed, $\mathcal{D}$-bimodule functor structures on $M_{g,h}(X)$ for $X \in \mathcal{C}_h$ are in bijection with central structures on X.

For every $g \in G$ choose a simple object $X_g \in \mathcal{C}_g$. Then $A_g := X_g \otimes X_g^*$ is an algebra in $\mathcal{D}$. The category of left A_g-modules in $\mathcal{C}$ is equivalent to $\mathcal{C}$ as a right $\mathcal{C}$-module category and the category of A_g-modules in $\mathcal{D}$ is equivalent to $\mathcal{C}_g$ as a right $\mathcal{D}$-module category.

It follows that for all $g, h \in G$ there is an equivalence $Y \mapsto X_g \otimes Y \otimes X_{hg}^*$ between $\mathcal{C}$ and the category of $A_g - A_{hg}$ bimodules in $\mathcal{C}$.

It restricts to an equivalence between $\mathcal{C}_h$ and the category of $A_g - A_{hg}$ bimodules in $\mathcal{D}$. It is easy to see that the latter equivalence coincides with (21).

The proof of equivalence (20) is completely similar. □

Let us define tensor functors

$$T_{g,h} := L^{-1}_{g,ghg^{-1}} R_{g,h} : \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_h) \to \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_{ghg^{-1}}), \quad g,h \in G, \tag{22}$$

and set

$$T_g := \bigoplus_{h\in G} T_{g,h} : \mathcal{Z}_{\mathcal{D}}(\mathcal{C}) \to \mathcal{Z}_{\mathcal{D}}(\mathcal{C}). \tag{23}$$

It follows that there is a natural family of isomorphisms:

$$c_{X,Y} : X \otimes Y \xrightarrow{\sim} T_g(Y) \otimes X, \qquad X \in \mathcal{C}_g,\, Y \in \mathcal{Z}_{\mathcal{D}}(\mathcal{C}), \quad g \in G, \tag{24}$$

satisfying natural compatibility conditions. Since the grading (18) is faithful we have $T_g(\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_h)) \subset \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_{ghg^{-1}})$.

Take $X_1 \in \mathcal{C}_{g_1}$, $X_2 \in \mathcal{C}_{g_2}$ and set $X = X_1 \otimes X_2$ in (24). We obtain a natural isomorphism

$$T_{g_1}T_{g_2}(Y) \otimes X_1 \otimes X_2 \xrightarrow{\sim} T_{g_1g_2}(Y) \otimes X_1 \otimes X_2.$$

and, hence, an isomorphism of functors $T_{g_1}T_{g_2} \xrightarrow{\sim} T_{g_1g_2}$. Thus, the assignment $g \mapsto T_g$ is an action of G on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ by tensor autoequivalences.

Suppose that X is an object in $\mathcal{Z}(\mathcal{C}_g)$. Then both sides of (24) have structure of objects in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ obtained by composing central structures of X and Y.

Lemma 3.2. *Isomorphisms* (24) *define a G-braiding on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$.*

Proof. That isomorphisms (24) are indeed morphisms in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ follows from commutativity of the diagram

(25)



where $(X, \gamma) \in \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g)$, $(Y, \delta) \in \mathcal{Z}_{\mathcal{D}}(\mathcal{C})$, and $V \in \mathcal{D}$. Indeed, the parallelogram in the middle commutes by naturality of c, and the two triangles commute since the natural isomorphisms $? \otimes Y \xrightarrow{\sim} T_g(Y) \otimes ? : \mathcal{C}_g \to \mathcal{C}_{gh}$, $g, h \in G$, commute with left and right actions of $\mathcal{D}$.

It is straightforward to check that isomorphisms $c_{X,Y}$ satisfy the compatibility conditions of Definition 2.10. □

The above constructions and arguments prove the following

Theorem 3.3. *Let G be a finite group and let $\mathcal{C}$ be a fusion category with a faithful G-grading whose trivial component is $\mathcal{D}$. The relative center $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ has a canonical structure of a braided G-crossed category.*

Remark 3.4. In particular, to every G-extension of a fusion category $\mathcal{D}$ we assigned an action of G by braided autoequivalences of $\mathcal{Z}(\mathcal{D})$. This assignment is studied in detail in [ENO3].

3.2. **The center $\mathcal{Z}(\mathcal{C})$ as an equivariantization.** As before, let G be a finite group and let $\mathcal{C}$ be a fusion category with a faithful G-grading (17). Let $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ be the braided G-crossed category constructed in Section 3.1.

Theorem 3.5. *There is an equivalence of braided fusion categories*

$$\mathcal{Z}_{\mathcal{D}}(\mathcal{C})^G \xrightarrow{\sim} \mathcal{Z}(\mathcal{C}). \tag{26}$$

Proof. We see from (24) that a G-equivariant object in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ has a structure of a central object in $\mathcal{C}$ defined as in (13). It follows from definitions that the corresponding tensor functor $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})^G \to \mathcal{Z}(\mathcal{C})$ is braided.

Conversely, given an object Y in $\mathcal{Z}(\mathcal{C})$ consider its forgetful image $\tilde{Y}$ in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. Combining the central structure of Y with isomorphism (24) we obtain natural isomorphisms

$$\tilde{Y} \otimes X \xrightarrow{\sim} T_g(\tilde{Y}) \otimes X, \qquad X \in \mathcal{C}_g,\ g \in G,$$

which give rise to a G-equivariant structure on $\tilde{Y}$. Hence, we have a tensor functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{Z}_{\mathcal{D}}(\mathcal{C})^G$. It is clear that the above two functors are quasi-inverses of each other. □

Let us describe the Tannakian subcategory $\mathcal{E} \cong \mathrm{Rep}(G) \subset \mathcal{Z}(\mathcal{C})$ corresponding to equivalence (26). For any representation $\pi : G \to GL(V)$ of the grading group G consider an object I_π in $\mathcal{Z}(\mathcal{C})$ where $I_\pi = V \otimes \mathbf{1}$ as an object of $\mathcal{C}$ with the permutation isomorphism

$$c_{I_\pi, X} := \pi(g) \otimes \mathrm{id}_X : I_\pi \otimes X \cong X \otimes I_\pi, \qquad \text{when } X \in \mathcal{C}_g. \tag{27}$$

Then $\mathcal{E}$ is the subcategory of $\mathcal{Z}(\mathcal{C})$ consisting of objects I_π, where π runs through all finite-dimensional representations of G.

Remark 3.6. Here is another description of the subcategory $\mathcal{E}$: it consists of all objects in $\mathcal{Z}(\mathcal{C})$ sent to Vec by the forgetful functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{Z}_{\mathcal{D}}(\mathcal{C})$.

Corollary 3.7. *Let $\mathcal{C}$ be a faithfully G-graded fusion category with the trivial component $\mathcal{D}$. Let $\mathcal{E} = \mathrm{Rep}(G) \subset \mathcal{Z}(\mathcal{C})$ be the Tannakian subcategory constructed above. Then the de-equivariantization category $(\mathcal{E}')_G$ is braided tensor equivalent to $\mathcal{Z}(\mathcal{D})$.*

Proof. The statement follows from Theorem 3.5 since $(\mathcal{E}')_G$ is the trivial component of the grading of $\mathcal{Z}(\mathcal{C})_G = \mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. □

Remark 3.8. The above assignment

$$\{G\text{-extensions of } \mathcal{D}\} \mapsto \{\text{braided } G\text{-crossed extensions of } \mathcal{Z}(\mathcal{D})\} \tag{28}$$

can be thought of as an analogue of the center construction for G-extensions.

Next, we describe simple objects of $\mathcal{Z}(\mathcal{C})$. For any conjugacy class K in G fix a representative $a_K \in K$. Let G_K denote the centralizer of a_K in G. Note that the action (23) of G on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ restricts to the action of G_K on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_{a_K})$.

Proposition 3.9. *There is a bijection between the set of isomorphism classes of simple objects of $\mathcal{Z}(\mathcal{C})$ and pairs (K, X), where K is a conjugacy class of G and X is a simple G_K-equivariant object of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_{a_K})$.*

Proof. By Theorem 3.5 we have $\mathcal{Z}(\mathcal{C}) \simeq \mathcal{Z}_{\mathcal{D}}(\mathcal{C})^G$ so the stated parameterization is immediate from the description of simple objects of the equivariantization category given in Proposition 2.7. □

3.3. **A criterion for a graded fusion category to be group-theoretical.** We have seen in Corollary 3.7 that $\mathcal{Z}(\mathcal{C})$ contains a Tannakian subcategory $\mathcal{E} = \mathrm{Rep}(G)$ such that the de-equivariantization $(\mathcal{E}')_G$ is braided equivalent to $\mathcal{Z}(\mathcal{D})$, where $\mathcal{D}$ is the trivial component of $\mathcal{C}$. Furthermore, by Remark 2.11, there is a canonical action of G on $\mathcal{Z}(\mathcal{D})$, by braided autoequivalences. By [DGNO], Tannakian subcategories of $\mathcal{Z}(\mathcal{C})$ containing $\mathcal{E}$ bijectively correspond to G-stable Tannakian subcategories of $(\mathcal{E}')_G \simeq \mathcal{Z}(\mathcal{D})$. Combining this observation with Theorem 2.5(ii) we obtain the following criterion.

Corollary 3.10. *A graded fusion category $\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g$, $\mathcal{C}_e = \mathcal{D}$, is group-theoretical if and only if $\mathcal{Z}(\mathcal{D})$ contains a G-stable Lagrangian subcategory.*

We will use Corollary 3.10 in Section 4.4 to characterize group-theoretical Tambara-Yamagami categories.

We can specialize Corollary 3.10 to equivariantization categories. Let G be a finite group acting on a fusion category $\mathcal{C}$. The equivariantization $\mathcal{C}^G$ is Morita equivalent to the crossed product category $\mathcal{C} \rtimes G$, see Section 2.4, therefore, $\mathcal{Z}(\mathcal{C}^G) \cong \mathcal{Z}(\mathcal{C} \rtimes G)$. Clearly, the trivial component of $\mathcal{Z}(\mathcal{C} \rtimes G)_G$ is $\mathcal{Z}(\mathcal{C})$ and the canonical action of G on $\mathcal{Z}(\mathcal{C})$ is induced from the action of G on $\mathcal{C}$ in an obvious way.

Corollary 3.11. *The equivariantization $\mathcal{C}^G$ is group-theoretical if and only if there exists a G-stable Lagrangian subcategory of $\mathcal{Z}(\mathcal{C})$.*

Remark 3.12. Let G act on $\mathcal{C}$ as before. One can check (independently from the results of this section) that the G-set of Lagrangian subcategories of $\mathcal{Z}(\mathcal{C})$ is isomorphic to the G-set of indecomposable pointed $\mathcal{C}$-module categories. This isomorphism is given by the map constructed in [NN, Theorem 4.17]. Thus, the criterion in Corollary 3.11 is the same as [Ni, Corollary 3.6].

3.4. **Example: the relative center of a crossed product category.** Let G be a finite group and let $g \mapsto T_g$, $g \in G$, be an action of G on a fusion category $\mathcal{D}$. Let $\mathcal{C} := \mathcal{D} \rtimes G$ be the crossed product category defined in Section 2.4. It has a natural grading

$$\mathcal{C} = \bigoplus_{g\in G} \mathcal{C}_g, \quad \text{where } \mathcal{C}_g = \{Y \boxtimes g \mid Y \in \mathcal{D}\}.$$

Let us describe the braided G-crossed fusion category structure on the relative center

$$\mathcal{Z}_{\mathcal{D}}(\mathcal{C}) = \bigoplus_{g\in G} \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g).$$

By definition, the objects of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g)$ are pairs $(Y \boxtimes g, \gamma)$, where $Y \in \mathcal{D}$ and

$$\gamma = \{\gamma_X : X \otimes Y \xrightarrow{\sim} Y \otimes T_g(X)\}_{X\in\mathcal{D}} \tag{29}$$

is a natural family of isomorphisms satisfying natural compatibility conditions. Thus, $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g)$ can be viewed as a "deformation" of $\mathcal{Z}(\mathcal{D})$ by means of T_g.

The action of G on $\mathcal{D}$ induces an action $h \mapsto \tilde{T}_h$ on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ defined as follows. Applying T_h, $h \in G$, to $\gamma_{T_{h^{-1}}(X)}$ in (29) we obtain an isomorphism

$$\tilde{\gamma}_X : X \otimes T_h(Y) \xrightarrow{\sim} T_h(Y) \otimes T_{hgh^{-1}}(X). \tag{30}$$

Set $\tilde{T}_h(Y \boxtimes g, \gamma) := (T_h(Y) \boxtimes hgh^{-1}, \tilde{\gamma})$. Thus, $\tilde{T}_h$ maps $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g)$ to $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_{hgh^{-1}})$.

Finally, the G-braiding between objects $(X \boxtimes h) \in \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_h)$ and $(Y \boxtimes g) \in \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_g)$ comes from the following isomorphism

$$\begin{aligned}(X \boxtimes h) \otimes (Y \boxtimes g) &= (X \otimes T_h(Y)) \boxtimes hg \\ &\xrightarrow{\tilde{\gamma}} (T_h(Y) \otimes T_{hgh^{-1}}(X)) \boxtimes hg \\ &= (T_h(Y) \boxtimes hgh^{-1}) \otimes (X \boxtimes h) \\ &= \tilde{T}_h(Y \boxtimes g) \otimes (X \boxtimes h).\end{aligned}$$

By Theorem 3.5, the category $\mathcal{Z}(\mathcal{D} \rtimes G) \cong \mathcal{Z}(\mathcal{D}^G)$ is equivalent to the equivariantization of the above braided G-crossed category.

4. The centers of Tambara-Yamagami categories

Our goal in this section is to apply techniques developed in Section 3 to Tambara-Yamagami categories introduced in [TY] (see Section 4.1 below for the definition). Namely, using the techniques in Section 3 we establish a criterion for a Tambara-Yamagami category to be group-theoretical. We then use this criterion together with Corollary 3.11 to produce a series of non group-theoretical semisimple Hopf algebras. In this section we assume that our ground field k is the field of complex numbers $\mathbb{C}$. We begin by recalling the definition of Tambara-Yamagami category.

4.1. **Definition of the Tambara-Yamagami category.** In [TY] D. Tambara and S. Yamagami completely classified all $\mathbb{Z}/2\mathbb{Z}$-graded fusion categories in which all but one simple object are invertible. They showed that any such category $\mathcal{TY}(A, \chi, \tau)$ is determined, up to an equivalence, by a finite Abelian group A, a non-degenerate symmetric bilinear form $\chi : A \times A \to k^\times$, and a square root $\tau \in k$ of $|A|^{-1}$. The category $\mathcal{TY}(A, \chi, \tau)$ is described as follows. It is a skeletal category (i.e., such that any two isomorphic objects are equal) with simple objects $\{a \mid a \in A\}$ and m, and tensor product

$$a \otimes b = a + b, \quad a \otimes m = m, \quad m \otimes a = m, \quad m \otimes m = \bigoplus_{a \in A} a,$$

for all $a, b \in A$, and the unit object $0 \in A$. The associativity constraints are given by

$$\begin{aligned}
\alpha_{a,b,c} &= \mathrm{id}_{a+b+c}, & \alpha_{a,b,m} &= \mathrm{id}_m, \\
\alpha_{a,m,b} &= \chi(a,b)\ \mathrm{id}_m, & \alpha_{m,a,b} &= \mathrm{id}_m, \\
\alpha_{a,m,m} &= \bigoplus_{b \in A} \mathrm{id}_b, & \alpha_{m,a,m} &= \bigoplus_{b \in A} \chi(a,b)\ \mathrm{id}_b, \\
\alpha_{m,m,a} &= \bigoplus_{b \in A} \mathrm{id}_b, & \alpha_{m,m,m} &= \bigoplus_{a,b \in A} \tau\chi(a,b)^{-1}\ \mathrm{id}_m\,.
\end{aligned}$$

The unit constraints are the identity maps. The category $\mathcal{TY}(A, \chi, \tau)$ is rigid with $a^* = -a$ and $m^* = m$ (with obvious evaluation and coevaluation maps).

Let $n := |A|$. The dimensions of simple objects of $\mathcal{TY}(A, \chi, \tau)$ are $\mathrm{FPdim}(a) = 1$, $a \in A$, and $\mathrm{FPdim}(m) = \sqrt{n}$. We have $\mathrm{FPdim}(\mathcal{TY}(A, \chi, \tau)) = 2n$.

Let $\mathbb{Z}/2\mathbb{Z} = \{1, \delta\}$. The $\mathbb{Z}/2\mathbb{Z}-$grading on $\mathcal{TY}(A, \chi, \tau)$ is

$$\mathcal{TY}(A, \chi, \tau) = \mathcal{TY}(A, \chi, \tau)_1 \oplus \mathcal{TY}(A, \chi, \tau)_\delta$$

where $\mathcal{TY}(A, \chi, \tau)_1$ is the full fusion subcategory generated by the invertible objects $a \in A$ and $\mathcal{TY}(A, \chi, \tau)_\delta$ is the full abelian subcategory generated by the object m.

Let $\mathcal{C} := \mathcal{TY}(A, \chi, \tau)$ and $\mathcal{D} := \mathcal{TY}(A, \chi, \tau)_1$.

4.2. **Braided $\mathbb{Z}/2\mathbb{Z}$-crossed category $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$.** First, let us describe the simple objects of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}) = \mathcal{Z}(\mathcal{C}_1) \oplus \mathcal{Z}_{\mathcal{D}}(\mathcal{C}_\delta)$. Let $\widehat{A} := \mathrm{Hom}(A, k^\times)$. Clearly, $\mathcal{Z}(\mathcal{C}_1) = \mathcal{Z}(\mathrm{Vec}_A)$, so its simple objects are parameterized by $(a, \phi) \in A \times \widehat{A}$. The object $X_{(a,\phi)}$ corresponding to such a pair is equal to a as an object of $\mathcal{C}$ and its central structure is given by

$$\phi(x)\,\mathrm{id}_{a+x} : x \otimes X_{(a,\phi)} \xrightarrow{\sim} X_{(a,\phi)} \otimes x. \tag{31}$$

Using Definition 2.1 we see that simple objects of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C}_\delta)$ are parameterized by functions $\rho : A \to k^\times$ satisfying

$$\rho(a+b) = \chi(a,b)^{-1}\rho(a)\rho(b), \qquad a, b \in A \tag{32}$$

(clearly, such functions form a torsor over $\widehat{A}$). The corresponding object Z_ρ is equal to m as an object of $\mathcal{C}$ and has the relative central structure

$$\rho(x)\,\mathrm{id}_m : x \otimes Z_\rho \xrightarrow{\sim} Z_\rho \otimes x, \quad x \in A. \tag{33}$$

Let $A \to \widehat{A} : a \mapsto \widehat{a}$ be the homomorphism defined by $\widehat{a}(x) = \chi(x,\, a)$. Similarly, let $\widehat{A} \to A : \phi \mapsto \widehat{\phi}$ be the homomorphism defined by $\phi(x) = \chi(x,\, \widehat{\phi})$ (recall that χ is non-degenerate). Clearly, these two maps are inverses of each other.

The fusion rules of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ are computed using formula (3) :

$$\begin{aligned} X_{(a,\phi)} \otimes X_{(b,\psi)} &= X_{(a+b,\phi+\psi)}, \\ X_{(a,\phi)} \otimes Z_\rho &= Z_{\rho\phi(-\widehat{a})}, \\ Z_\rho \otimes X_{(a,\phi)} &= Z_{\rho\phi(-\widehat{a})}, \\ Z_{\rho'} \otimes Z_\rho &= \bigoplus_{a \in A} X_{(a, \widehat{a}\rho'/\overline{\rho})}. \end{aligned}$$

We have $X^*_{(a,\phi)} = X_{(-a,-\phi)}$ and $Z^*_\rho = Z_{\overline{\rho}}$, where $\overline{\rho}(x) = \rho(-x)$, $x \in A$.

Using the construction given in Section 3.1 we see that the action of $\mathbb{Z}/2\mathbb{Z}$ on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ is given by

$$T_1 = \mathrm{id}_{\mathcal{Z}_{\mathcal{D}}(\mathcal{C})}; \qquad T_\delta(X_{(a,\phi)}) = X_{(-\widehat{\phi},\, -\widehat{a})}, \quad T_\delta(Z_\rho) = Z_\rho. \tag{34}$$

The monoidal functor structure on $\mathbb{Z}/2\mathbb{Z} \to \mathrm{Aut}_\otimes(\mathcal{Z}_{\mathcal{D}}(\mathcal{C}))$ is given by the natural isomorphism $\gamma := \gamma_{\delta,\delta} : T_\delta \circ T_\delta \xrightarrow{\sim} T_1$ defined by

$$\gamma_{X_{(a,\phi)}} = \phi(a)\,\mathrm{id}_{X_{(a,\phi)}}, \qquad \gamma_{Z_\rho} = \left(\tau \sum_{x \in A} \rho(x)^{-1}\right) \mathrm{id}_{Z_\rho}.$$

The crossed braiding morphisms on $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ are given by

$$\begin{aligned} c_{X_{(a,\phi)}, X_{(b,\psi)}} &= \psi(a)\,\mathrm{id}_{a+b} : X_{(a,\phi)} \otimes X_{(b,\psi)} \xrightarrow{\sim} X_{(b,\psi)} \otimes X_{(a,\phi)} \\ c_{X_{(a,\phi)}, Z_\rho} &= \rho(a)\,\mathrm{id}_m : X_{(a,\phi)} \otimes Z_\rho \xrightarrow{\sim} Z_\rho \otimes X_{(a,\phi)} \\ c_{Z_\rho, X_{(a,\phi)}} &= \mathrm{id}_m : Z_\rho \otimes X_{(a,\phi)} \xrightarrow{\sim} X_{(-\widehat{\phi},\, -\widehat{a})} \otimes Z_\rho \\ c_{Z_{\rho'}, Z_\rho} &= \oplus_{a \in A}\ \rho(-a)^{-1}\,\mathrm{id}_a : Z_{\rho'} \otimes Z_\rho \xrightarrow{\sim} Z_\rho \otimes Z_{\rho'}. \end{aligned}$$

4.3. **The equivariantization category $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})^{\mathbb{Z}/2\mathbb{Z}}$.** A simple calculation of $\mathbb{Z}/2\mathbb{Z}$-equivariant objects in $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ establishes the following.

Proposition 4.1. *The following is a complete list of simple objects of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})^{\mathbb{Z}/2\mathbb{Z}} \cong \mathcal{Z}(\mathcal{TY}(A,\chi,\tau))$ up to an isomorphism:*

(1) *$2n$ invertible objects parameterized by pairs (a,ϵ), where $a \in A$ and $\epsilon^2 = \chi(a,a)^{-1}$. The corresponding object $X_{a,\epsilon}$ is equal to $X_{(a,-\widehat{a})}$ as an object of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ and has $\mathbb{Z}/2\mathbb{Z}-$equivariant structure*

$$\epsilon\,\mathrm{id}_{X_{(a,-\widehat{a})}} : T_\delta(X_{(a,-\widehat{a})}) \xrightarrow{\sim} X_{(a,-\widehat{a})};$$

(2) *$\frac{n(n-1)}{2}$ two-dimensional objects parameterized by unordered pairs (a,b) of distinct objects in A. The corresponding object $Y_{a,b}$ is equal to $X_{(a,-\widehat{b})} \oplus X_{(b,-\widehat{a})}$ as an object of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ and has $\mathbb{Z}/2\mathbb{Z}-$equivariant structure*

$$\left(\mathrm{id}_{X_{(a,-\widehat{b})}} \oplus \chi(a,b)^{-1}\,\mathrm{id}_{X_{(b,-\widehat{a})}}\right) : T_\delta(X_{(a,-\widehat{b})} \oplus X_{(b,-\widehat{a})}) \xrightarrow{\sim} X_{(a,-\widehat{b})} \oplus X_{(b,-\widehat{a})};$$

(3) *$2n$ $\sqrt{n}-$dimensional objects parameterized by pairs (ρ,Δ), where $\rho : A \to k^\times$ satisfies (32) and $\Delta^2 = \tau \sum_{x\in A} \rho(x)^{-1}$. The corresponding object $Z_{\rho,\Delta}$ is equal to Z_ρ as an object of $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ and has $\mathbb{Z}/2\mathbb{Z}-$equivariant structure*

$$\Delta\,\mathrm{id}_{Z_\rho} : T_\delta(Z_\rho) \xrightarrow{\sim} Z_\rho.$$

Recall from [ENO1] that in a braided fusion category of an integer Frobenius-Perron dimension there is a canonical choice of a twist θ such that the categorical dimensions of objects coincide with their Frobenius-Perron dimensions. Namely, for any simple object X the scalar θ_X is defined in such a way that the composition

$$\mathbf{1} \xrightarrow{\mathrm{coev}_X} X \otimes X^* \xrightarrow{\theta_X c_{X,X^*}} X^* \otimes X \xrightarrow{\mathrm{ev}_X} \mathbf{1} \tag{35}$$

is equal to $\mathrm{FPdim}(X)\,\mathrm{id}_X$.

Let θ be the canonical twist on $\mathcal{Z}(\mathcal{C})$. Using the above observation, explicit formulas from Subsection 4.2, and Section 2.6, we immediately obtain the following.

$$\theta_{X_{a,\epsilon}} = \chi(a,a)^{-1}, \qquad \theta_{Y_{a,b}} = \chi(a,b)^{-1}, \qquad \theta_{Z_{\rho,\Delta}} = \Delta.$$

Using the fusion rules of $\mathcal{Z}(\mathcal{C})$ (which may be computed using the explicit formulas in Subsection 4.2), values of the twists above, and the well known formula

$$S_{X,Y} = \theta_X^{-1}\theta_Y^{-1} \sum_Z N_{X,Y}^Z \theta_Z d_Z \tag{36}$$

we obtain the S- and T-matrices of $\mathcal{Z}(\mathcal{C})$:

$$S_{X_{a,\epsilon},X_{a',\epsilon'}} = \chi(a,a')^2, \qquad S_{X_{a,\epsilon},Y_{b,c}} = 2\chi(a,b+c),$$
$$S_{X_{a,\epsilon},Z_{\rho,\Delta}} = \epsilon\sqrt{n}\rho(a), \qquad S_{Y_{a,b},Y_{c,d}} = 2\left(\chi(a,d)\chi(b,c) + \chi(a,c)\chi(b,d)\right),$$
$$S_{Y_{a,b},Z_{\rho,\Delta}} = 0, \qquad S_{Z_{\rho,\Delta},Z_{\rho',\Delta'}} = \frac{1}{\Delta\Delta'}\sum_{a\in A} \chi(a,a)^2\rho(a)\rho'(a).$$

$$T_{X_{a,\epsilon}} = \chi(a,a)^{-1}, \quad T_{Y_{a,b}} = \chi(a,b)^{-1}, \quad T_{Z_{\rho,\Delta}} = \Delta.$$

Proposition 4.2. *The maximal pointed subcategory of $\mathcal{Z}(\mathcal{C})$ is non-degenerate if and only if $|A|$ is odd.*

Proof. Let $a \in A$ be an element of order 2. Then $X_{a,\epsilon}$ centralizes every invertible object of $\mathcal{Z}(\mathcal{C})$. □

Remark 4.3. We note that simple objects and the S- and T-matrices of $\mathcal{Z}(\mathcal{C})$ were described by Izumi in [I] using very different methods.

4.4. **A criterion for a Tambara-Yamagami category to be group-theoretical.** The group $A \times \widehat{A}$ is equipped with a canonical non-degenerate quadratic form $q : A \times \widehat{A} \to k^\times$ given by

$$q((a, \phi)) := \phi(a), \quad (a, \phi) \in A \times \widehat{A}.$$

We will call a subgroup $B \subset A \times \widehat{A}$ *Lagrangian* if $q|_B = 1$ and $B = B^\perp$ with respect to the bilinear form defined by q. Lagrangian subgroups of $A \times \widehat{A}$ correspond to Lagrangian subcategories of $\mathcal{Z}(\mathrm{Vec}_A) \cong \mathrm{Vec}_{A \times \widehat{A}}$.

The braided tensor autoequivalence T_δ of $\mathcal{Z}(\mathrm{Vec}_A)$ defined in Section 4.2 determines an order 2 automorphism of $A \times \widehat{A}$, which we denote simply by δ:

$$\delta((a, \phi)) = (-\widehat{\phi}, -\widehat{a}), \quad (a, \phi) \in A \times \widehat{A}. \tag{37}$$

Definition 4.4. We will say that a subgroup $L \subset A$ is *Lagrangian (with respect to χ)* if $L = L^\perp$ with respect to the inner product on A given by χ. Equivalently, $|L|^2 = |A|$ and $\chi|_L = 1$.

Lemma 4.5. *Let A be an Abelian 2-group such that $|A| = 2^{2n}$ and let χ be a non-degenerate symmetric bilinear form on A. Then A contains a Lagrangian subgroup.*

Proof. It suffices to show that A contains an isotropic element, i.e., an element $x \in A$, $x \neq 0$, such that $\chi(x, x) = 1$. Then one can pass from A to $\langle x \rangle^\perp / \langle x \rangle$ and use induction.

Suppose that A is cyclic with a generator a. Then $2^{2n}a = 0$ and $\chi(a, a)$ is a 2^{2n}-th root of unity, hence $\chi(2^n a, 2^n a) = \chi(a, a)^{2^{2n}} = 1$.

If A is not cyclic then it contains a subgroup $A_0 = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$. Let x_1, x_2 be distinct non-zero elements of A_0. Suppose $\chi(x_i, x_i) \neq 1$, $i = 1, 2$. Then $\chi(x_i, x_i) = -1$ and $\chi(x_1 + x_2, x_1 + x_2) = 1$, as desired. □

Theorem 4.6. *Let $\mathcal{C} = \mathcal{TY}(A, \chi, \tau)$ be a Tambara-Yamagami fusion category. Then $\mathcal{C}$ is group-theoretical if and only if A contains a Lagrangian subgroup (with respect to χ).*

Proof. By Corollary 3.10, $\mathcal{C}$ is group-theoretical if and only if $\mathcal{Z}(\mathcal{D})$ contains a T_δ-stable Lagrangian subcategory. Equivalently, $\mathcal{C}$ is group-theoretical if and only if $A \times \widehat{A}$ contains a Lagrangian subgroup B stable under the action

$$(a, \phi) \mapsto (\widehat{\phi}, \widehat{a}). \tag{38}$$

This condition on B is the same as being stable under the action of δ from (37).

Let L be a Lagrangian (with respect to χ) subgroup of A and let $\widehat{L} := \{\widehat{a} \mid a \in L\}$. Then $L \times \widehat{L}$ is a Lagrangian subgroup of $A \times \widehat{A}$ stable under (38). Hence $\mathcal{C}$ is group-theoretical.

Conversely, suppose that $\mathcal{C}$ is group-theoretical. Let us write $A = A_{\mathrm{even}} \oplus A_{\mathrm{odd}}$, where A_{even} is the Sylow 2-subgroup of A and A_{odd} is the maximal odd order subgroup of A. Since $|A|$ must be a square, we conclude that $|A_{\mathrm{even}}|$ is a square, and so A_{even} contains a Lagrangian subgroup with respect to $\chi|_{A_{\mathrm{even}}}$ by Lemma 4.5.

So it remains to show that A_{odd} contains a Lagrangian subgroup with respect to $\chi|_{A_{\mathrm{odd}}}$. For this end we may assume that $|A|$ is odd. Let $B \subset A \times \widehat{A}$ be a

Lagrangian subgroup stable under (38). Then $B = B_+ \oplus B_-$, where

$$B_\pm := \{(a, \pm\widehat{a}) \mid (a, \pm\widehat{a}) \in B\}.$$

Let $L_\pm = B_\pm \cap (A \times \{1\})$. Then $|L_+||L_-| = |A|$, and $\chi|_{L_\pm} = 1$. Hence, $L_\pm$ are Lagrangian subgroups of A. □

Remark 4.7. It was observed in [ENO1, Remark 8.48] that for an odd prime p and elliptic bicharacter χ on $A = (\mathbb{Z}/p\mathbb{Z})^2$ the category $\mathcal{TY}((\mathbb{Z}/p\mathbb{Z})^2, \chi, \tau)$ is not group-theoretical. The criterion from Theorem 4.6 extends this observation.

4.5. **A series of non group-theoretical semisimple Hopf algebras obtained from Tambara-Yamagami categories.** Here we apply Corollary 3.11 to produce a series of non group-theoretical fusion categories admitting fiber functors (i.e., representation categories of non group-theoretical semisimple Hopf algebras), generalizing examples constructed in [Ni].

Let A be a finite Abelian group with a non-degenerate bilinear form χ. Let $\mathrm{Aut}(A, \chi)$ denote the group of automorphisms of A preserving χ.

The following proposition was proved in [Ni, Proposition 2.10].

Proposition 4.8. *There is an action of* $\mathrm{Aut}(A, \chi)$ *on* $\mathcal{TY}(A, \chi, \tau)$ *given by* $g \mapsto T_g$, *where*

$$T_g(A) = g(a), \quad T_g(m) = m, \qquad a \in A,\ g \in \mathrm{Aut}(A, \chi),$$

with the tensor structure of T_g *given by identity morphisms.*

Corollary 4.9. *Let* G *be a subgroup of* $\mathrm{Aut}(A, \chi)$. *Then the fusion category* $\mathcal{TY}(A, \chi, \tau)^G$ *is group-theoretical if and only if there is a Lagrangian subgroup of* (A, χ) *stable under the action of* G.

Proof. Combine Corollary 3.11 and Theorem 4.6. □

We will say that a non-degenerate symmetric bilinear form $\chi : A \times A \to k^\times$ is *hyperbolic* if there are Lagrangian subgroups $L, L' \subset A$ such that $A = L \oplus L'$. Note that in this case L' is isomorphic to the group $\widehat{L} = \mathrm{Hom}(L, k^\times)$ of characters of L and χ is identified with the canonical bilinear form on $L \oplus \widehat{L}$.

It was shown by D. Tambara in [Ta1] that when $n = |A|$ is odd the category $\mathcal{TY}(A, \chi, \tau)$ admits a fiber functor (i.e., $\mathcal{TY}(A, \chi, \tau)$ is equivalent to the representation category of a semisimple Hopf algebra) if and only if τ^{-1} is a positive integer and χ is hyperbolic.

Corollary 4.10. *Let* p *be an odd prime, let* $L = (\mathbb{Z}/p\mathbb{Z})^N$, $N \geq 1$, *let* $A = L \oplus \widehat{L}$, *and let* $\chi : A \times A \to k^\times$ *be the canonical bilinear form defined by*

$$\chi((a, \phi), (b, \psi)) = \psi(a)\phi(b), \quad a, b \in A,\ \phi, \psi \in \widehat{A}.$$

Suppose that G *is a subgroup of* $\mathrm{Aut}(A, \chi)$ *not contained in any conjugate of* $\mathrm{Aut}(L) \subset \mathrm{Aut}(A, \chi)$. *Then the equivariantization category* $\mathcal{TY}(A, \chi, p^{-N})^G$ *is a non group-theoretical fusion category equivalent to the representation category of a semisimple Hopf algebra of dimension* $2p^{2N}|G|$.

Proof. Note that $\mathrm{Aut}(A, \chi)$ acts transitively on the set of Lagrangian subgroups of (A, χ) and the stabilizer of L is $\mathrm{Aut}(L)$. Apply Corollary 4.9. □

Remark 4.11. The series of fusion categories in Corollary 4.10 extends the one constructed in [Ni], where the case of $N = 1$ and $G = \mathbb{Z}/2\mathbb{Z}$ was considered.

5. Examples of modular categories arising from quadratic forms

As before, let $\mathcal{C} := \mathcal{TY}(A, \chi, \tau)$ be a Tambara-Yamagami category and let $\mathcal{D} := \mathcal{TY}(A, \chi, \tau)_1$ be the trivial component of $\mathbb{Z}/2\mathbb{Z}-$grading of $\mathcal{TY}(A, \chi, \tau)$. In this section we assume that our ground field k is the field of complex numbers $\mathbb{C}$.

Suppose that the symmetric bicharacter $\chi : A \times A \to k^\times$ comes from a quadratic form on A, i.e., there is a function $q : A \to k^\times$ such that

$$q(a+b) = q(a)q(b)\chi(a,b), \quad a, b \in A \quad \text{and} \quad q(-a) = q(a).$$

¿From the description obtained in Section 4.2 we observe that $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$ contains a fusion subcategory spanned by the simple objects $X_{(a,\widehat{a})}, a \in A$, and $Z_{q^{-1}}$. It is clear from the Tambara-Yamagami classification in Section 4.1 that this category is equivalent to $\mathcal{C}$.

Proposition 5.1. *Suppose that the symmetric bicharacter χ comes from a quadratic form on A. Then $\mathcal{C}$ admits a $\mathbb{Z}/2\mathbb{Z}$-crossed braided category structure. The equivariantization $\mathcal{C}^{\mathbb{Z}/2\mathbb{Z}}$ is non-degenerate if and only if $|A|$ is odd.*

Proof. Clearly, $\mathcal{C}$ inherits the $\mathbb{Z}/2\mathbb{Z}$-crossed braided category structure from $\mathcal{Z}_{\mathcal{D}}(\mathcal{C})$. The non-degeneracy claim follows from Proposition 4.2 and Remark 2.13. □

Let us assume that $n := |A|$ is odd. Then χ corresponds to a unique quadratic form q. Let $\mathcal{E}(q, \pm) := \mathcal{C}^{\mathbb{Z}/2\mathbb{Z}}$ be the modular category constructed in Proposition 5.1 (the $\pm$ corresponding to $\tau = \pm\frac{1}{\sqrt{n}}$, respectively). In what follows we describe the fusion rules and S- and T-matrices of $\mathcal{E}(q, \pm)$.

5.1. **Fusion rules of $\mathcal{E}$.** Clearly, $\mathcal{E}(q, \pm)$ is a fusion category of dimension $4n$. It has the following simple objects:

two invertible objects, $\mathbf{1} = X_+$ and X_-,
$\frac{n-1}{2}$ two-dimensional objects Y_a, $a \in A - \{0\}$ (with $Y_{-a} = Y_a$)
two $\sqrt{n}$-dimensional objects Z_l, $l \in \mathbb{Z}/2\mathbb{Z}$.

Here we simplify the notation used in Subsection 4.3 and denote

$$X_\pm := X_{0,\pm 1},\ Y_a := Y_{a,-a}, \text{ and } Z_l := Z_{q^{-1},\Delta_l},$$

where $\Delta_l, l \in \mathbb{Z}/2\mathbb{Z}$, are distinct square roots of $\pm\frac{1}{\sqrt{n}}\sum_{a\in A} q(a)$.

The fusion rules of $\mathcal{E}(q, \pm)$ are given by:

$$\begin{aligned}
&X_- \otimes X_- = X_+, && X_\pm \otimes Y_a = Y_a, && X_+ \otimes Z_l &= Z_l,\\
&X_- \otimes Z_l &= Z_{l+1}, && Y_a \otimes Y_b &= Y_{a+b} \oplus Y_{a-b}, && Y_a \otimes Y_a &= X_+ \oplus X_- \oplus Y_{2a},\\
&Y_a \otimes Z_l &= Z_0 \oplus Z_1, && Z_l \otimes Z_l &= X_+ \oplus (\oplus Y_a), && Z_l \otimes Z_{l+1} &= X_- \oplus (\oplus Y_a),
\end{aligned}$$

where $a, b \in A$ $(a \neq b)$ and $l \in \mathbb{Z}/2\mathbb{Z}$. All objects of $\mathcal{E}(q, \pm)$ are self-dual.

Remark 5.2. Note that the fusion rules of $\mathcal{E}(q, \pm)$ do not depend on the quadratic form q and the number τ. We show below that the S- and T-matrices of $\mathcal{E}(q, \pm)$ do depend on q and τ.

5.2. **S- and T-matrices of $\mathcal{E}$.**

Lemma 5.3. *The Gauss sums corresponding to q and q^2 are equal up to a sign, i.e.,*

$$\frac{\sum_{a\in A} q(a)^2}{\sum_{a\in A} q(a)} \in \{\pm 1\}.$$

Proof. Consider the group $A \times A$ with a non-degenerate quadratic form $Q = q \times q$. The Gaussian sum for this form is

$$\tau(A \times A, Q) = \sum_{a,b \in A} q(a)q(b) = \tau(A, q)^2.$$

The restriction of Q on the diagonal subgroup $D := \{(a, a) \mid a \in A\}$ is non-degenerate since $|A|$ is odd. The restriction of Q on the orthogonal complement $D^\perp = \{(a, -a) \mid a \in A\}$ is non-degenerate as well. By the multiplicativity of Gaussian sums we have

$$\tau(A \times A, Q) = \tau(D, Q)\tau(D^\perp, Q) = (\sum_{a \in A} q(a)^2)^2,$$

which implies the result. □

Using the formulas for the S- and T- matrices of $\mathcal{Z}(\mathcal{C})$ given in Subsection 4.3 we can write down the S- and T- matrices of $\mathcal{E}(q, \pm)$:

$$S_{X_\pm, X_\pm} = 1, \qquad S_{X_\mp, X_\pm} = 1, \qquad S_{X_\pm, Y_a} = 2,$$

$$S_{X_+, Z_l} = \sqrt{n}, \qquad S_{X_-, Z_l} = -\sqrt{n}, \qquad S_{Y_a, Y_b} = 2\left(\frac{q(a+b)^2}{q(a)^2 q(b)^2} + \frac{q(a)^2 q(b)^2}{q(a+b)^2}\right),$$

$$S_{Y_a, Z_l} = 0, \qquad S_{Z_l, Z_l} = \begin{cases} \pm\sqrt{n}, & \text{if the Gauss sums of } q \text{ and } q^2 \text{ coincide,} \\ \mp\sqrt{n}, & \text{otherwise,} \end{cases}$$

$$S_{Z_l, Z_{l+1}} = \begin{cases} \mp\sqrt{n}, & \text{if the Gauss sums of } q \text{ and } q^2 \text{ coincide,} \\ \pm\sqrt{n}, & \text{otherwise.} \end{cases}$$

$$T_{X_\pm} = 1, \quad T_{Y_a} = q(a)^2, \quad T_{Z_l} = \Delta_l.$$

(Recall that $\Delta_l, l \in \mathbb{Z}/2\mathbb{Z}$, are distinct square roots of $\pm\frac{1}{\sqrt{n}} \sum_{a \in A} q(a)$.)

5.3. **Example with $A = \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$.** Let p be an odd prime and let $A := \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$. Let $\left(\frac{\cdot}{p}\right)$ denote the Legendre symbol modulo p, i.e., $\left(\frac{a}{p}\right) = 1$ if $a \in (\mathbb{Z}/p\mathbb{Z})^\times$ is a square modulo p and -1 otherwise.

Let $a, b \in (\mathbb{Z}/p\mathbb{Z})^\times$ and $\xi := e^{\frac{2\pi i}{p}}$. Consider the following nondegenerate quadratic form q on A:

$$q(x_1, x_2) = \xi^{ax_1^2 - bx_2^2}.$$

It is hyperbolic if $\left(\frac{ab}{p}\right) = 1$ and elliptic if $\left(\frac{ab}{p}\right) = -1$.

We will need the following.

Lemma 5.4. *For every $a, b \in A^\times$, we have*

$$\sum_{x \in \mathbb{Z}/p\mathbb{Z}} \xi^{ax^2} = \begin{cases} \left(\frac{a}{p}\right)\sqrt{p}, & \text{if } p \equiv 1 \,(\mathrm{mod}\, 4), \\ \left(\frac{a}{p}\right) i\sqrt{p}, & \text{if } p \equiv 3 \,(\mathrm{mod}\, 4) \end{cases}$$

and

$$\sum_{(x_1, x_2) \in \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}} \xi^{ax_1^2 - bx_2^2} = \left(\frac{ab}{p}\right) p.$$

Proof. The first assertion is well known, see for example [R]. The second assertion is an easy consequence of the first. □

Using Lemma 5.4 we can explicitly write the S-matrix of $\mathcal{E}(q,\pm)$:

$$\begin{aligned} S_{X_\pm,X_\pm} &= 1, & S_{X_\mp,X_\pm} &= 1, & S_{X_\pm,Y_{(x_1,x_2)}} &= 2,\\ S_{X_+,Z_l} &= p, & S_{X_-,Z_l} &= -p, & S_{Y_{(x_1,x_2)},Y_{(y_1,y_2)}} &= 4\,\mathrm{Re}(\xi^{4ax_1y_1-4bx_2y_2}),\\ S_{Y_{(x_1,x_2)},Z_l} &= 0, & S_{Z_l,Z_l} &= \pm p, & S_{Z_l,Z_{l+1}} &= \mp p, \end{aligned}$$

and its T-matrix:

$$T_{X_\pm} = 1, \quad T_{Y_(x_1,x_2)} = \xi^{2ax_1^2-2bx_2^2}, \quad T_{Z_l} = \Delta_l,$$

where $\Delta_l, l \in \mathbb{Z}/2\mathbb{Z}$, are distinct square roots of $\pm\left(\frac{ab}{p}\right)$.

The central charge of the modular category $\mathcal{E}(q,\pm)$ is

$$\zeta(\mathcal{E}(q,\pm)) = \left(\frac{ab}{p}\right).$$

Below we give the S- and T-matrices of the modular category $\mathcal{E}(q,\pm)$ for $p=3$. Order simple objects of $\mathcal{E}(q,\pm)$ as follows: $\mathbf{1}, X_-, Y_{(0,1)}, Y_{(1,0)}, Y_{(1,1)}, Y_{(1,2)}, Z_+, Z_-$. There are four modular categories $\mathcal{E}(q,\pm)$ of dimension 36 corresponding to the choices of hyperbolic/elliptic q and $\tau = \pm\frac{1}{3}$.

(a) When q is hyperbolic we have:

$$S = \begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 2 & 3 & 3 \\ 1 & 1 & 2 & 2 & 2 & 2 & -3 & 3 \\ 2 & 2 & -2 & 4 & -2 & -2 & 0 & 0 \\ 2 & 2 & 4 & -2 & -2 & -2 & 0 & 0 \\ 2 & 2 & -2 & -2 & 4 & -2 & 0 & 0 \\ 2 & 2 & -2 & -2 & -2 & 4 & 0 & 0 \\ 3 & -3 & 0 & 0 & 0 & 0 & \pm 3 & \mp 3 \\ 3 & -3 & 0 & 0 & 0 & 0 & \mp 3 & \pm 3, \end{pmatrix}$$

$$T = \mathrm{diag}\{1, 1, \xi^2, \xi, 1, 1, 1, -1\} \quad \text{when } \tau = \frac{1}{3},$$

$$T = \mathrm{diag}\{1, 1, \xi^2, \xi, 1, 1, i, -i\} \quad \text{when } \tau = -\frac{1}{3}.$$

Note that both the corresponding modular categories are group-theoretical with central charge 1; in fact the one with $\tau = \frac{1}{3}$ is equivalent to the representation category of the double $D(S_3)$ of the symmetric group S_3 and the one with $\tau = -\frac{1}{3}$ is equivalent to the twisted double of S_3.

(b) When q is elliptic we have:

$$S = \begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 2 & 3 & 3 \\ 1 & 1 & 2 & 2 & 2 & 2 & -3 & 3 \\ 2 & 2 & -2 & 4 & -2 & -2 & 0 & 0 \\ 2 & 2 & 4 & -2 & -2 & -2 & 0 & 0 \\ 2 & 2 & -2 & -2 & -2 & 4 & 0 & 0 \\ 2 & 2 & -2 & -2 & 4 & -2 & 0 & 0 \\ 3 & -3 & 0 & 0 & 0 & 0 & \pm 3 & \mp 3 \\ 3 & -3 & 0 & 0 & 0 & 0 & \mp 3 & \pm 3, \end{pmatrix}$$

$$T = \operatorname{diag}\{1, 1, \xi, \xi, \xi^2, \xi^2, i, -i\} \quad \text{when } \tau = \frac{1}{3},$$

$$T = \operatorname{diag}\{1, 1, \xi, \xi, \xi^2, \xi^2, 1, -1\} \quad \text{when } \tau = -\frac{1}{3}.$$

Both the corresponding modular categories are not group-theoretical. They both have central charge -1 and so are not equivalent to centers of fusion categories. In particular, they are not equivalent to representation categories of any twisted group doubles.

5.4. **Example with $A = \mathbb{Z}/p\mathbb{Z}$.** Let p be an odd prime and let $A := \mathbb{Z}/p\mathbb{Z}$. Let $a \in (\mathbb{Z}/p\mathbb{Z})^\times$ and $\xi := e^{\frac{2\pi i}{p}}$. Upto isomorphism the are two nondegenerate quadratic forms q on A:

$$q(x) = \xi^{ax^2},$$

one corresponding to $\left(\frac{a}{p}\right) = 1$ and another to $\left(\frac{a}{p}\right) = -1$.

Using Lemma 5.4 we can explicitly write the S-matrix of $\mathcal{E}(q, \pm)$:

$$\begin{aligned} &S_{X_\pm, X_\pm} = 1, && S_{X_\mp, X_\pm} = 1, && S_{X_\pm, Y_x} = 2, \\ &S_{X_+, Z_l} = \sqrt{p}, && S_{X_-, Z_l} = -\sqrt{p}, && S_{Y_x, Y_y} = 4\,\mathrm{Re}(\xi^{4axy}), \\ &S_{Y_a, Z_l} = 0, && S_{Z_l, Z_l} = \pm\left(\frac{2}{p}\right)\sqrt{p}, && S_{Z_l, Z_{l+1}} = \mp\left(\frac{2}{p}\right)\sqrt{p}. \end{aligned}$$

$$T_{X_\pm} = 1, \quad T_{Y_x} = \xi^{-2ax^2}, \quad T_{Z_l} = \Delta_l,$$

where

$$\Delta_l, l \in \mathbb{Z}/2\mathbb{Z}, \text{ are distinct} \begin{cases} \text{square roots of } \pm\left(\frac{a}{p}\right), \text{ if } p \equiv 1 \pmod 4, \\ \text{square roots of } \pm\left(\frac{a}{p}\right) i, \text{ if } p \equiv 3 \pmod 4. \end{cases}$$

The central charge of the modular category $\mathcal{E}(q, \pm)$ is

$$\zeta(\mathcal{E}(q, \pm)) = \begin{cases} \left(\frac{2a}{p}\right), \text{ if } p \equiv 1 \pmod 4, \\ -\left(\frac{2a}{p}\right) i, \text{ if } p \equiv 3 \pmod 4. \end{cases}$$

Below we give the S- and T-matrices of the modular category $\mathcal{E}(q, \pm)$ for $p = 3$ and 5. For $p = 3$ we order the simple objects as $\mathbf{1}, X_-, Y_1, Z_0, Z_1$ and for $p = 5$ we order them as $\mathbf{1}, X_-, Y_1, Y_2, Z_0, Z_1$. (In (c) and (d) below, $\xi = e^{\frac{2\pi i}{5}}$.)

(a) When $p = 3$ and $a = 1$ we have:

$$S = \begin{pmatrix} 1 & 1 & 2 & \sqrt{3} & \sqrt{3} \\ 1 & 1 & 2 & -\sqrt{3} & -\sqrt{3} \\ 2 & 2 & -2 & 0 & 0 \\ \sqrt{3} & -\sqrt{3} & 0 & \mp\sqrt{3} & \pm\sqrt{3} \\ \sqrt{3} & -\sqrt{3} & 0 & \pm\sqrt{3} & \mp\sqrt{3} \end{pmatrix}$$

$$T = \operatorname{diag}\left\{1, 1, \frac{-1+i\sqrt{3}}{2}, \frac{1+i}{\sqrt{2}}, \frac{-1-i}{\sqrt{2}}\right\} \quad \text{when } \tau = \frac{1}{\sqrt{3}},$$

$$T = \operatorname{diag}\left\{1, 1, \frac{-1+i\sqrt{3}}{2}, \frac{1-i}{\sqrt{2}}, \frac{-1+i}{\sqrt{2}}\right\} \quad \text{when } \tau = -\frac{1}{\sqrt{3}}.$$

The central charge of both the corresponding modular categories is i.

(b) When $p = 3$ and $a = 2$ we have:

$$S = \text{ the } S\text{-matrix in (a)}$$

$$T = \operatorname{diag}\left\{1, 1, \frac{-1-i\sqrt{3}}{2}, \frac{1-i}{\sqrt{2}}, \frac{-1+i}{\sqrt{2}}\right\} \quad \text{when } \tau = \frac{1}{\sqrt{3}},$$

$$T = \operatorname{diag}\left\{1, 1, \frac{-1-i\sqrt{3}}{2}, \frac{1+i}{\sqrt{2}}, \frac{-1-i}{\sqrt{2}}\right\} \quad \text{when } \tau = \frac{1}{\sqrt{3}}.$$

The central charge of both the corresponding modular categories is $-i$.

(c) When $p = 5$ and $a = 1$ we have:

$$S = \begin{pmatrix} 1 & 1 & 2 & 2 & \sqrt{5} & \sqrt{5} \\ 1 & 1 & 2 & 2 & -\sqrt{5} & -\sqrt{5} \\ 2 & 2 & \sqrt{5}-1 & -\sqrt{5}-1 & 0 & 0 \\ 2 & 2 & -\sqrt{5}-1 & \sqrt{5}-1 & 0 & 0 \\ \sqrt{5} & -\sqrt{5} & 0 & 0 & \mp\sqrt{5} & \pm\sqrt{5} \\ \sqrt{5} & -\sqrt{5} & 0 & 0 & \pm\sqrt{5} & \mp\sqrt{5} \end{pmatrix}$$

$$T = \operatorname{diag}\left\{1, 1, \xi^3, \xi^2, 1, -1\right\} \quad \text{when } \tau = \frac{1}{\sqrt{5}},$$

$$T = \operatorname{diag}\left\{1, 1, \xi^3, \xi^2, i, -i\right\} \quad \text{when } \tau = -\frac{1}{\sqrt{5}}.$$

The central charge of both the corresponding modular categories is -1.

(d) When $p = 5$ and $a = 2$ we have:

$$S = \begin{pmatrix} 1 & 1 & 2 & 2 & \sqrt{5} & \sqrt{5} \\ 1 & 1 & 2 & 2 & -\sqrt{5} & -\sqrt{5} \\ 2 & 2 & -\sqrt{5}-1 & \sqrt{5}-1 & 0 & 0 \\ 2 & 2 & \sqrt{5}-1 & -\sqrt{5}-1 & 0 & 0 \\ \sqrt{5} & -\sqrt{5} & 0 & 0 & \mp\sqrt{5} & \pm\sqrt{5} \\ \sqrt{5} & -\sqrt{5} & 0 & 0 & \pm\sqrt{5} & \mp\sqrt{5} \end{pmatrix}$$

$$T = \operatorname{diag}\left\{1, 1, \xi, \xi^4, i, -i\right\} \quad \text{when } \tau = \frac{1}{\sqrt{5}},$$

$$T = \operatorname{diag}\left\{1, 1, \xi, \xi^4, 1, -1\right\} \quad \text{when } \tau = -\frac{1}{\sqrt{5}}.$$

The central charge of both the corresponding modular categories is 1.

6. Appendix: Zeroes in S-matrices

There is a classical result of Burnside in character theory saying that if χ is an irreducible character of a finite group G and $\chi(1) > 1$ then $\chi(g) = 0$ for some $g \in G$, see [BZ, Chapter 21].

In this appendix we establish a categorical analogue of this result for weakly integral modular categories. Recall [ENO2] that a fusion category $\mathcal{C}$ is called *weakly integral* if its Frobenius-Perron dimension is an integer. In this case the Frobenius-Perron dimension of every simple object of $\mathcal{C}$ is the square root of an integer [ENO1].

Let $\mathcal{C}$ be a weakly integral modular category with the S-matrix S. Let $\mathcal{O}(\mathcal{C})$ denote the set of all (representatives of isomorphism classes of) simple object of $\mathcal{C}$. Given $X \in \mathcal{O}(\mathcal{C})$ define the following sets:

$$T_X = \{Y \in \mathcal{O}(\mathcal{C}) \mid S_{X,Y} = 0\},$$
$$D_X = \mathcal{O}(\mathcal{C}) - (T_X \cup \{\mathbf{1}\}).$$

Clearly, we have a partition $\mathcal{O}(\mathcal{C}) = T_X \cup D_X \cup \{\mathbf{1}\}$. Let $\mathcal{T}_X$ and $\mathcal{D}_X$ be full Abelian subcategories of $\mathcal{C}$ generated by T_X and D_X, respectively.

Let K be the field extension of $\mathbb{Q}$ generated by the entries of S. It is known [dBG, CG] that there is a root of unity ξ such that $K \subset \mathbb{Q}(\xi)$. In particular, the operation of taking the square of an absolute value of an element of S is well defined. Let $G := \operatorname{Gal}(K/\mathbb{Q})$. Every element $\sigma \in G$ comes from a permutation σ of $\mathcal{O}(\mathcal{C})$ such that $\sigma(S_{X,Y}) = S_{X,\sigma(Y)}$ for all $X, Y \in \mathcal{O}(\mathcal{C})$.

Let $\mathcal{C}$ be a weakly integral modular category. It was shown in [ENO1] that there is a canonical spherical structure on $\mathcal{C}$ such that categorical dimensions in $\mathcal{C}$ coincide with Frobenius-Perron dimensions. Let us fix this structure for the reminder of this section. For any $X \in \mathcal{O}(\mathcal{C})$ let d_X denote the dimension of X. For any full abelian subcategory $\mathcal{A}$ of $\mathcal{C}$ let $\dim(\mathcal{A})$ denote the sum of squares of dimensions of simple objects of $\mathcal{A}$.

Theorem 6.1. *Let $\mathcal{C}$ be a weakly integral modular category with the S-matrix S. Then T_X is not empty for every non-invertible simple object X of $\mathcal{C}$. That is, every row (column) of S corresponding to a non-invertible simple object contains at least one zero entry.*

Proof. Note that the statement of Proposition does not depend on the choice of spherical structure.

We have $\sum_{Y\in\mathcal{O}(\mathcal{C})}|S_{X,Y}|^2=\dim(\mathcal{C})$, hence,

(39)
$$1=\frac{\dim(\mathcal{C})}{d_X^2}-\sum_{Y\in D_X}\left|\frac{S_{X,Y}}{d_X}\right|^2=\frac{1+\dim(\mathcal{T}_X)}{d_X^2}-\left(\sum_{Y\in D_X}\left|\frac{S_{X,Y}}{d_X}\right|^2-\frac{\dim(\mathcal{D}_X)}{d_X^2}\right),$$

where d_X denotes the dimension of X. It suffices to check that

$$\frac{1}{\dim(\mathcal{D}_X)}\sum_{Y\in D_X}\left|\frac{S_{X,Y}}{d_X}\right|^2\geq\frac{1}{d_X^2} \tag{40}$$

since then (39) implies that $1\leq\frac{1+\dim(\mathcal{T}_X)}{d_X^2}$, whence

$$\dim(\mathcal{T}_X)\geq d_X^2-1. \tag{41}$$

But X is non-invertible so $d_X>1$ and $\mathcal{T}_X\neq 0$.

Rewriting the left hand side of (40) as the sum of $\dim(\mathcal{D}_X)$ terms and using the inequality of arithmetic and geometric means we obtain

$$\begin{aligned}\frac{1}{\dim(\mathcal{D}_X)}\sum_{Y\in D_X}\left|\frac{S_{X,Y}}{d_X}\right|^2&=\frac{1}{\dim(\mathcal{D}_X)}\sum_{Y\in D_X}d_Y^2\left|\frac{S_{X,Y}}{d_Xd_Y}\right|^2\\&\geq\frac{1}{d_X^2}\left(\prod_{Y\in D_X}\left|\frac{S_{X,Y}}{d_Y}\right|^{2d_Y^2}\right)^{\frac{1}{\dim(D_X)}}.\end{aligned}$$

The set D_X is clearly stable under all automorphisms in the Galois group, and hence so is the product $\prod_{Y\in D_X}\left|\frac{S_{X,Y}}{d_Y}\right|^{2d_Y^2}$. Therefore, this product belongs to $\mathbb{Q}$. Its factors are squares of absolute values of characters of $K_0(\mathcal{C})$ on X and hence are algebraic integers. Since all factors are positive, the product is ≥ 1, which implies (40). □

For $X\in\mathcal{O}(\mathcal{C})$ define

$$U_X=\{Y\in\mathcal{O}(\mathcal{C})\mid|S_{X,Y}|=d_Y\}.$$

Let $\mathcal{U}_X$ be the full Abelian subcategory of $\mathcal{C}$ generated by U_X.

Proposition 6.2. *Let $\mathcal{C}$ be a weakly integral modular category and let X be a simple non-invertible object in $\mathcal{C}$. Then*

$$3\dim(\mathcal{T}_X)+\dim(\mathcal{U}_X)>\dim(\mathcal{C}). \tag{42}$$

Proof. We may assume $d_X\geq\sqrt{2}$.

We will use the following theorem of Siegel [Si] from number theory. Let $K/\mathbb{Q}$ be a finite Galois extension with the Galois group $G=\mathrm{Gal}(K/\mathbb{Q})$. Let α be a totally positive algebraic integer in K, $\alpha\neq 1$. Then

$$\frac{1}{|G|}\sum_{\sigma\in G}\sigma(\alpha)\geq\frac{3}{2}.$$

We apply this to the situation when K is the extension of $\mathbb{Q}$ generated by entries of S. We compute

$$\begin{aligned}\dim(\mathcal{C}) &= \sum_{Y\in\mathcal{C}} |S_{X,Y}|^2 \\ &= d_X^2 + \sum_{Y\in U_X} d_Y^2 + \sum_{Y\in\mathcal{O}(\mathcal{C})-(T_X\cup U_X\cup\{\mathbf{1}\})} |S_{X,Y}|^2 \\ &= d_X^2 + \dim(\mathcal{U}_X) + \sum_{Y\in\mathcal{O}(\mathcal{C})-(T_X\cup U_X\cup\{\mathbf{1}\})} d_Y^2 \left(\frac{1}{|G|}\sum_{\sigma\in G}\sigma\left(\frac{|S_{X,Y}|^2}{d_Y^2}\right)\right) \\ &\geq 2 + \dim(\mathcal{U}_X) + \frac{3}{2}(\dim(\mathcal{C}) - \dim(\mathcal{T}_X) - \dim(\mathcal{U}_X) - 1),\end{aligned}$$

therefore $3\dim(\mathcal{T}_X) + \dim(\mathcal{U}_X) \geq \dim(\mathcal{C}) + 1 > \dim(\mathcal{C})$, as required. □

Remark 6.3. Our proofs of Theorem 6.1 and Proposition 6.2 imitate the corresponding proofs for group characters given in [BZ].

References

[AG] S. Arkhipov and D. Gaitsgory, *Another realization of the category of modules over the small quantum group*, Adv. Math. **173** (2003), no. 1, 114-143.

[BK] B. Bakalov, A. Kirillov Jr., *Lectures on Tensor categories and modular functors*, AMS, (2001).

[Br] A. Bruguières, *Catégories prémodulaires, modularization et invariants des variétés de dimension* 3, Mathematische Annalen, **316** (2000), no. 2, 215-236.

[BZ] Ya.G. Berkovich, E.M. Zhmud', *Characters of Finite Groups. Part 2*, Translations of Mathematical Monographs, volume 181, American Mathematical Society.

[CG] A. Coste, T. Gannon, *Remarks on Galois symmetry in rational conformal field theories*, Phys. Lett. B **323** (1994), no. 3-4, 316-321.

[dBG] J. de Boere, J. Goeree, *Markov traces and II_1 factors in conformal field theory*, Comm. Math. Phys. **139** (1991), no. 2, 267-304.

[D] P. Deligne, *Catégories tannakiennes*, The Grothendieck Festschrift, Vol. II, 111–195, Progr. Math., 87, Birkhäuser Boston, Boston, MA, 1990.

[DGNO] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik, *On braided fusion categories I* (a preliminary version: *Group-theoretical properties of nilpotent modular categories*, arXiv:0704.0195v2 [math.QA]).

[ENO1] P. Etingof, D. Nikshych, V. Ostrik, *On fusion categories*, Annals of Mathematics **162** (2005), 581-642.

[ENO2] P. Etingof, D. Nikshych, V. Ostrik, *Weakly group-theoretical and solvable fusion categories*, arXiv:0809.3031.

[ENO3] P. Etingof, D. Nikshych, V. Ostrik, *Fusion categories and homotopy theory*, in preparation.

[G] D. Gaitsgory, *The notion of category over an algebraic stack*, arXiv:0507192v1 [math.AG].

[GN] S. Gelaki, D. Nikshych, *Nilpotent fusion categories*, Advances in Mathematics **217** (2008) 1053-1071.

[I] M. Izumi, *The structure of sectors associated with Longo-Rehren inclusions II. Examples*, Reviews in Mathematical Physics, **13**(2001), Issue 05, 603-674.

[K] C. Kassel, *Quantum groups*, Graduate Texts in Mathematics, **155**, Springer-Verlag, (1995).

[Ki] A. Kirillov, Jr. *Modular categories and orbifold models II*, arXiv:0110221 [math.QA].

[Ma] S. Majid, *Representations, duals and quantum doubles of monoidal categories*, Rend. Circ. Mat. Palermo (2) Suppl., **26** (1991), 197-206.

[Mu1] M. Müger, *Galois theory for braided tensor categories and the modular closure*, Adv. Math. **150** (2000), no. 2, 151-201.

[Mu2] M. Müger, On the structure of modular categories, *Proc. Lond. Math. Soc.*, **87** (2003), 291-308.

[Mu3] M. Müger, *From subfactors to categories and topology. I. Frobenius algebras in and Morita equivalence of tensor categories.*, J. Pure Appl. Algebra **180** (2003), no. 1-2, 81–157.

[Mu4] M. Müger, *Galois extensions of braided tensor categories and braided crossed G-categories*, J. Algebra **277** (2004), no. 1, 256–281.

[NN] D. Naidu and D. Nikshych, *Lagrangian subcategories and braided tensor equivalences of twisted quantum doubles of finite groups*, Communications in Mathematical Physics **279** (2008), no. 3, 845–872.

[Ni] D. Nikshych, *Non group-theoretical semisimple Hopf algebras from group actions on fusion categories*, Selecta Mathematica, **14** (2008) 145-161.

[O] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups, **8** (2003), 177-206.

[R] I. K. Rosen, *A classical introduction to modern number theory*, Graduate Texts in Mathematics, **84**, Springer-Verlag, (1990).

[S] J. A. Siehler, *Braided Near-group Categories*, arXiv: 0011037 [math.QA].

[Si] C.L. Siegel, *The trace of totally positive and real algebraic integers*, Ann. Math., 46 (1945), 302-312.

[Ta1] D. Tambara, *Representations of tensor categories with fusion rules of self-duality for abelian groups*, Isr. J. Math. **118** (2000), 29-60.

[Ta2] D. Tambara, *Invariants and semi-direct products for finite group actions on tensor categories*, J. Math. Soc. Japan **53** (2001), no. 2, 429-456.

[TY] D. Tambara, S. Yamagami, *Tensor categories with fusion rules of self-duality for finite abelian groups*, J. Algebra **209** (1998), no. 2, 692-707.

[Tu1] V. Turaev, *Homotopy field theory in dimension 3 and crossed group-categories*, arXiv:math/0005291.

[Tu2] V. Turaev, *Crossed group-categories*, Arabian Journal for Science and Engineering, **33**, no. 2C, 484-503 (2008).

S.G.: Department of Mathematics, Technion-Israel Institute of Technology, Haifa 32000, Israel.

E-mail address: gelaki@math.technion.ac.il

D.N.: Department of Mathematics, Texas A&M University, College Station, TX 77843, USA.

E-mail address: dnaidu@math.tamu.edu

D.N.: Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824, USA.

E-mail address: nikshych@math.unh.edu